

Ans.  
Q1

5

## 5-SUBSTITUTED ARYLPYRIMIDINES

This application claims the benefit of U.S. Provisional Application Serial No. 60/189,774 filed March 16, 2000 and of U.S. Provisional Application Serial No. 60/206,454 filed May 22, 2000, the teachings of which are incorporated herein by reference.

10

### FIELD OF THE INVENTION

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

The present invention relates to novel substituted arylpyrimidine compounds that bind with high selectivity and/ or high affinity to CRF1 receptors (Corticotropin Releasing Factor 1 Receptors). This invention also relates to pharmaceutical compositions comprising such compounds and to the use of such compounds in treatment of psychiatric disorders and neurological diseases, including major depression, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy and feeding disorders, as well as treatment of immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological disturbance and stress. Additionally this invention relates to the use such compounds as probes for the localization of CRF1 receptors in cells and tissues.

### BACKGROUND OF THE INVENTION

Corticotropin releasing factor (CRF), a 41 amino acid peptide, is the primary physiological regulator of proopiomelanocortin (POMC) derived peptide secretion from the anterior pituitary gland. In addition to its endocrine role at the pituitary gland, immunohistochemical localization of CRF has demonstrated that the hormone has a broad extrahypothalamic distribution in the central nervous system and produces a wide spectrum of autonomic, electrophysiological and behavioral effects consistent with a neurotransmitter or neuromodulator role in brain. There is also evidence that CRF plays a significant role in integrating the response of the immune system to physiological, psychological, and immunological stressors.

Clinical data provide evidence that CRF has a role in psychiatric disorders and neurological diseases including depression, anxiety-related disorders and feeding disorders. A role for CRF has also been postulated in the etiology and pathophysiology of Alzheimer's disease, Parkinson's disease, Huntington's disease, progressive supranuclear palsy and amyotrophic lateral sclerosis as they relate to the dysfunction of CRF neurons in the central nervous system.

In affective disorder, or major depression, the concentration of CRF is significantly increased in the cerebral spinal fluid (CSF) of drug-free individuals. Furthermore, the density of CRF receptors is significantly decreased in the frontal cortex of suicide victims, consistent with a hypersecretion of CRF. In addition, there is a blunted adrenocorticotropin (ACTH) response to CRF (i.v. administered) observed in depressed patients. Preclinical studies in rats and non-human primates provide additional support for the hypothesis that hypersecretion of CRF may be involved in the symptoms seen in human depression. There is also preliminary evidence that tricyclic antidepressants can alter CRF levels and thus modulate the numbers of CRF receptors in brain.

CRF has also been implicated in the etiology of anxiety-related disorders. CRF produces anxiogenic effects in animals and interactions between benzodiazepine / non-benzodiazepine anxiolytics and CRF have been demonstrated in a variety of behavioral anxiety models.

Preliminary studies using the putative CRF receptor antagonist  $\alpha$ -helical ovine CRF (9-41) in a variety of behavioral paradigms demonstrate that the antagonist produces "anxiolytic-like" effects that are qualitatively similar to the benzodiazepines. Neurochemical, endocrine and receptor binding studies have all demonstrated interactions between CRF and benzodiazepine anxiolytics providing further evidence for the involvement of CRF in these disorders. Chlordiazepoxide attenuates the "anxiogenic" effects of CRF in both the conflict test and in the acoustic startle test in rats. The benzodiazepine receptor antagonist Ro 15-1788, which was without behavioral activity alone in the operant conflict test, reversed the effects of CRF in a dose-dependent manner, while the benzodiazepine inverse agonist FG 7142 enhanced the actions of CRF.


CRF has also been implicated in the pathogenesis of certain immunological, cardiovascular or heart-related diseases such as hypertension, tachycardia and congestive heart failure, stroke and osteoporosis, as well as in premature birth, psychosocial dwarfism, stress-

induced fever, ulcer, diarrhea, post-operative ileus and colonic hypersensitivity associated with psychopathological disturbance and stress.

The mechanisms and sites of action through which conventional anxiolytics and antidepressants produce their therapeutic effects remain to be fully elucidated. It has been hypothesized however, that they are involved in the suppression of CRF hypersecretion that is observed in these disorders. Of particular interest are that preliminary studies examining the effects of a CRF receptor antagonist peptide ( $\alpha$ -helical CRF<sub>9-41</sub>) in a variety of behavioral paradigms have demonstrated that the CRF antagonist produces "anxiolytic-like" effects qualitatively similar to the benzodiazepines.

### Description of the Related Art

A number of small molecule compounds for the treatment of CRF related disorders have been disclosed in the literature [for a review see J. McCarthy et al. *Curr. Pharm. Des.* 5: 289(1999)]. McCarthy et al. (WO 96/39400) have disclosed aryl pyrimidine derivatives of the general formula




wherein X, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are defined therein, for use as CRF receptor in the treatment of central nervous system disorders. The McCarthy application discloses arylpyrimidine compounds that contain a disubstituted amino group (NR<sub>1</sub>R<sub>2</sub>) in the 4-position of the pyrimidine ring. It is therefore surprising that the novel pyrimidines of this invention, which lack the corresponding disubstituted NR<sub>1</sub>R<sub>2</sub> group in the 4-position of the pyrimidine ring, are also CRF receptor antagonists.

### SUMMARY OF THE INVENTION

The invention provides novel compounds of Formula I (shown below), and pharmaceutical compositions comprising compounds of Formula I and at least one pharmaceutically acceptable carrier or excipient. Such arylpyrimidines bind to cell surface

receptors, preferably G-coupled protein receptors, especially CRF receptors and most preferably CRF1 receptors. Preferred compounds of the invention exhibit high affinity for CRF 1 receptors. Additionally, preferred compounds of the invention also exhibit high specificity for CRF1 receptors.

5 Typically preferred compounds of the invention include those of Formula I:



Formula I

and the pharmaceutically acceptable salts thereof, wherein:

10 Ar is phenyl, 1- or 2-naphthyl, each of which is mono-, di-, or tri-substituted or mono-, di-, or tri-substituted heteroaryl having from about 5 to about 7 ring members and 1 to about 4 heteroatoms in the ring, the heteroatoms independently selected from the group consisting of N, O and S;

15 R<sub>1</sub> and R<sub>3</sub> are independently chosen from hydrogen, halogen, cyano, nitro, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted (cycloalkyl)alkyl, optionally substituted alkylthio, optionally substituted alkylsulfinyl, optionally substituted alkylsulfonyl, or optionally substituted mono- or dialkylcarboxamide, with the proviso that R<sub>1</sub> and R<sub>3</sub> are not both hydrogen; and

20 R<sub>2</sub> is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted aminoalkyl, optionally substituted mono or dialkylamino, optionally substituted alkylthio, optionally substituted alkylsulfinyl, optionally substituted alkylsulfonyl, optionally substituted mono or dialkylcarboxamide, optionally substituted carbocyclic aryl or optionally substituted heteroaryl having from 1 to 3 rings, and 3 to 8 ring members in each ring and 1 to about 3 heteroatoms.

25

Particular embodiments of this invention include compounds of Formula I in which R<sub>1</sub> and R<sub>3</sub> are as defined above for Formula I, Ar is phenyl which is mono-, di-, or tri-substituted; and

R<sub>2</sub> is selected from optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aminoalkyl, optionally substituted mono or dialkylamino, optionally substituted alkylthio, optionally substituted alkylsulfinyl, optionally substituted alkylsulfonyl, optionally substituted mono and dialkylcarboxamide, or

5 R<sub>2</sub> is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidinyl, pyridizynyl, and thiophenyl, each of which is optionally mono-, di-, or tri-substituted.

Particular embodiments of this invention also include compounds of Formula I in which R<sub>1</sub> and R<sub>3</sub> are as defined above for Formula I, Ar is mono-, di-, or trisubstituted phenyl; and

10 R<sub>2</sub> is selected from optionally substituted alkoxy, optionally substituted aminoalkyl, and optionally substituted mono or dialkylamino.

The invention further comprises methods of treating patients suffering from certain disorders with an effective amount of a compound of the invention. These disorders include  
15 CNS disorders, particularly affective disorders, anxiety disorders, stress-related disorders, eating disorders and substance abuse. The patient suffering from these disorders may be a human or other animal (preferably a mammal), such as a domesticated companion animal (pet) or a livestock animal.

According to yet another aspect, the present invention provides pharmaceutical  
20 compositions comprising compounds of Formula I or the pharmaceutically acceptable salts or solvates thereof, which compositions are useful for the treatment of the above-recited disorders. The invention further provides methods of treating patients suffering from any of the above-recited disorders with an effective amount of a compound or composition of the invention.

Additionally this invention relates to the use of the compounds of the invention  
25 (particularly labeled compounds of this invention) as probes for the localization of receptors in cells and tissues and as standards and reagents for use in determining the receptor-binding characteristics of test compounds.

Preferred arylpyrimidines of the invention exhibit good activity in standard *in vitro* receptor binding assays, specifically the assay as specified in Example 96, which follows and is  
30 defined below. Particularly preferred arylpyrimidines of the invention have an IC<sub>50</sub> of about 1 micromolar or less, still more preferably an IC<sub>50</sub> of about 100 nanomolar or less even more

preferably an IC<sub>50</sub> of about 10 nanomolar or less or even 1 nanomolar or less in such a defined standard *in vitro* CRF receptor binding assay as exemplified by Example 96 which follows.

### DETAILED DESCRIPTION OF THE INVENTION

In addition to compounds of Formula I, shown and described above, the invention also provides compounds and the pharmaceutically acceptable salts of Formula I, which will be referred to as compounds of **Formula Ia** in which:

R<sub>1</sub> and R<sub>3</sub> are independently selected from hydrogen, halogen, cyano, C<sub>1-6</sub> alkyl, (C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl, -O(C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl, halo(C<sub>1-6</sub>)alkyl, -O(halo(C<sub>1-6</sub>)alkyl), -O(C<sub>1-6</sub>alkyl), and S(O)<sub>n</sub>(C<sub>1-6</sub>alkyl),

where each alkyl is independently straight, branched, or cyclic, may contain 1 or more double or triple bonds, and is optionally substituted with one or more substituents independently chosen from halogen, hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino,

and

where each C<sub>3-7</sub>cycloalkyl is optionally substituted by one or more substituents independently chosen from halogen, hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino,

with the proviso that not both R<sub>1</sub> and R<sub>3</sub> are hydrogen;

R<sub>2</sub> is selected from the group consisting of -XR<sub>A</sub> and Y; and

Ar is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidinyl, pyridizynyl, and thiophenyl, each of which is mono-, di-, or tri-substituted with R<sub>C</sub>;

R<sub>A</sub> and R<sub>B</sub>, which may be the same or different, are independently selected at each occurrence from:

hydrogen and straight, branched, or cyclic alkyl groups, including (cycloalkyl)alkyl groups consisting of 1 to 8 carbon atoms, which straight, branched, or cyclic alkyl groups may contain one or more double or triple bonds, each of which 1 to 8 carbon atoms may be further substituted with one or more substituent(s) independently selected from oxo, hydroxy, halogen, cyano, amino, C<sub>1-6</sub>alkoxy, -NH(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -NHC(=O)(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)C(=O)(C<sub>1-6</sub>alkyl), -NHS(O)<sub>n</sub>(C<sub>1-6</sub>alkyl), -S(O)<sub>n</sub>(C<sub>1-6</sub>alkyl), -S(O)<sub>n</sub>NH(C<sub>1-6</sub>alkyl), -S(O)<sub>n</sub>N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), and Z;

R<sub>C</sub> is independently selected at each occurrence from halogen, cyano, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, hydroxy, amino, C<sub>1-6</sub>alkyl substituted with 0-2 R<sub>D</sub>, C<sub>2-6</sub> alkenyl substituted with 0-2 R<sub>D</sub>, C<sub>2-6</sub>alkynyl substituted with 0-2 R<sub>D</sub>, C<sub>3-7</sub>cycloalkyl substituted with 0-2 R<sub>D</sub>, (C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl substituted with 0-2 R<sub>D</sub>, C<sub>1-6</sub>alkoxy substituted with 0-2 R<sub>D</sub>, -NH(C<sub>1-6</sub>alkyl) substituted with 0-2 R<sub>D</sub>, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl) each C<sub>1-6</sub>alkyl independently substituted with 0-2 R<sub>D</sub>, -XR<sub>A</sub>, and Y;

R<sub>D</sub> is independently selected at each occurrence from the group consisting of halogen, hydroxy, cyano, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), -S(O)<sub>n</sub>(alkyl), halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, CO(C<sub>1-4</sub>alkyl), CONH(C<sub>1-4</sub>alkyl), CON(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), -XR<sub>A</sub>, and Y;

X is independently selected at each occurrence from the group consisting of -CH<sub>2</sub>-, -CHR<sub>B</sub>-, -O-, -C(=O)-, -C(=O)O-, -S(O)<sub>n</sub>-, -NH-, -NR<sub>B</sub>-, -C(=O)NH-, -C(=O)NR<sub>B</sub>-, -S(O)<sub>n</sub>NH-, -S(O)<sub>n</sub>NR<sub>B</sub>-, -OC(=S)S-, -NHC(=O)-, -NR<sub>B</sub>C(=O)-, -NHS(O)<sub>n</sub>-, -OSiH<sub>n</sub>(C<sub>1-4</sub>alkyl)<sub>2-n</sub>-, and -NR<sub>B</sub>S(O)<sub>n</sub>-;

Y and Z are independently selected at each occurrence from: 3- to 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, amino, cyano, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and -S(O)<sub>n</sub>(alkyl),

said 3- to 7-membered heterocyclic groups containing one or more heteroatom(s) independently selected from N, O, and S, with the point of attachment being either carbon or nitrogen; and

n is independently selected at each occurrence from 0, 1, and 2.

Particular embodiments of the invention include compounds and salts of Formula Ia in which Ar is phenyl, mono-, di-, or tri-substituted with R<sub>C</sub> and R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> are as defined for Formula Ia.


The invention further includes compounds and salts of Formula Ia in which Ar is phenyl mono-, di-, or tri-substituted with R<sub>C</sub>; R<sub>1</sub> and R<sub>3</sub> are independently selected groups (1) halogen and (2) C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, (C<sub>3-7</sub>cycloalkyl)C<sub>1-3</sub>alkyl, (C<sub>3-7</sub>cycloalkyl) C<sub>1-3</sub>alkoxy, where each

member of group (2) is unsubstituted or substituted by 1-3 groups independently chosen from hydroxy, amino, cyano, and halogen.

Also provided by the invention are compounds and salts of Formula Ia in which Ar is phenyl mono-, di-, or tri-substituted with  $R_C$ ; and  $R_A$  and  $R_B$ , which may be the same or different, are independently selected at each occurrence from straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which alkyl groups may contain one or more double or triple bonds.


Further provided as embodiments of the invention are compounds and salts of Formula Ia in which Ar is phenyl mono-, di-, or tri-substituted with  $R_C$ ;  $R_A$  and  $R_B$ , which may be the same or different, are independently selected at each occurrence from straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which alkyl groups may contain one or more double or triple bonds; and  $R_1$  and  $R_3$  are independently selected from groups (1) halogen and (2)  $C_{1-3}$ alkyl,  $C_{1-3}$ alkoxy,  $(C_{3-7}$ cycloalkyl) $C_{1-3}$ alkyl,  $(C_{3-7}$ cycloalkyl)  $C_{1-3}$ alkoxy, where each member of group (2) is unsubstituted or substituted by 1-3 groups independently chosen from hydroxy, amino, cyano, and halogen.

Also provided as embodiments of the invention are compounds and salts of Formula Ia having the formula



wherein  $Ar$ ,  $R_1$  and  $R_3$  are as defined for Formula Ia, and  $R_X$  and  $R_Y$  are the same or different and are independently selected from hydrogen and  $C_1 - C_6$  alkyl; or

$NR_XR_Y$  represents Formula Ib:




**Formula Ib**



wherein z is 0 or 1; and W is CR<sub>A</sub>R<sub>B</sub>, NR<sub>B</sub>, or O.

A preferred embodiment of the invention includes compounds of Formula A



Formula A

and the pharmaceutically acceptable salt thereof, wherein:

R<sub>X</sub> and R<sub>Y</sub> are the same or different and are independently selected from:

- a) hydrogen,
- b) -(C=O)alkyl<sub>A</sub>, wherein alkyl<sub>A</sub> is a straight or branched alkyl group having from 1 to 8 carbon atoms;
- c) straight, branched, or cyclic alkyl groups, including cycloalkyl(alkyl)groups, said alkyl groups having from 1 to 8 carbon atoms and optionally containing one or more double or triple bonds, each of which alkyl groups may be further substituted with one or more substituent(s) independently selected from:
  - i) hydroxy, halogen, amino, cyano, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -NH(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and
  - ii) 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents independently selected from halogen, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and -S(O)<sub>n</sub>(alkyl), wherein said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) independently selected from N, O, and S, with the point of attachment being either carbon or nitrogen,

R<sub>1</sub> and R<sub>3</sub> are independently selected from hydrogen, halogen, cyano, C<sub>1-6</sub> alkyl<sub>1</sub>, (C<sub>3-7</sub>cycloalkyl<sub>1</sub>)C<sub>1-4</sub>alkyl<sub>1</sub>, -O(C<sub>3-7</sub>cycloalkyl<sub>1</sub>)C<sub>1-4</sub>alkyl<sub>1</sub>, halo(C<sub>1-6</sub>)alkyl<sub>1</sub>, -O(halo(C<sub>1-6</sub>)alkyl<sub>1</sub>), -O(C<sub>1-6</sub>alkyl<sub>1</sub>), and S(O)<sub>n</sub>(C<sub>1-6</sub>alkyl<sub>1</sub>),

where each said alkyl<sub>1</sub> is straight, branched, or cyclic and may contain 1 or more double or triple bonds, and is optionally substituted by one or more substituents independently chosen from halogen, hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino,

and

where said C<sub>3-7</sub>cycloalkyl<sub>1</sub> is optionally substituted by one or more substituents independently chosen from halogen, hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino

with the proviso that not both R<sub>1</sub> and R<sub>3</sub> are hydrogen; and

Ar is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidinyl, and

thiophenyl, each of which is mono-, di-, or tri-substituted with R<sub>C</sub>;

R<sub>A</sub> and R<sub>B</sub>, which may be the same or different, are independently selected at each occurrence from the group consisting of:

hydrogen and straight, branched, or cyclic alkyl groups, including (cycloalkyl)alkyl

groups, consisting of 1 to 8 carbon atoms, which may contain one or more double or

triple bonds, each of which may be further substituted with one or more substituent(s)

independently selected from oxo, hydroxy, halogen, nitro, cyano, C<sub>1-6</sub>alkoxy, -NH(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -NHC(=O)(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)C(=O)(C<sub>1-6</sub>alkyl), -

NHS(O)<sub>n</sub>(C<sub>1-6</sub>alkyl), -S(O)<sub>n</sub>(C<sub>1-6</sub>alkyl), -S(O)<sub>n</sub>NH(C<sub>1-6</sub>alkyl), -S(O)<sub>n</sub>N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), and Z;

R<sub>C</sub> is independently selected at each occurrence from halogen, cyano, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, hydroxy, amino, and C<sub>1-6</sub>alkyl substituted with 0-2 R<sub>D</sub>, C<sub>2-6</sub>alkenyl substituted

with 0-2 R<sub>D</sub>, C<sub>2-6</sub>alkynyl substituted with 0-2 R<sub>D</sub>, C<sub>3-7</sub>cycloalkyl substituted with 0-2 R<sub>D</sub>, (C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl substituted with 0-2 R<sub>D</sub>, C<sub>1-6</sub>alkoxy substituted with 0-2 R<sub>D</sub>, -

NH(C<sub>1-6</sub>alkyl) substituted with 0-2 R<sub>D</sub>, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl) each C<sub>1-4</sub>alkyl

independently substituted with 0-2 R<sub>D</sub>, -XR<sub>A</sub>, and Y, with the proviso that at least one of

the positions ortho or para to the point of attachment of Ar to the pyrimidine ring shown in Formula A is substituted;

R<sub>D</sub> is independently selected at each occurrence the group consisting of halogen, hydroxy, cyano,

C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), -S(O)<sub>n</sub>(alkyl) halo(C<sub>1-</sub>

$_4$ alkyl, halo( $C_{1-4}$ alkoxy, CO( $C_{1-4}$ alkyl), CONH( $C_{1-4}$ alkyl), CON( $C_{1-4}$ alkyl)( $C_{1-4}$ alkyl), -XR<sub>A</sub>, and Y;

X is independently selected at each occurrence from the group consisting of -CH<sub>2</sub>-, -CHR<sub>B</sub>-, -O-, -C(=O)-, -C(=O)O-, -S(O)<sub>n</sub>-, -NH-, -NR<sub>B</sub>-, -C(=O)NH-, -C(=O)NR<sub>B</sub>-, -S(O)<sub>n</sub>NH-, -S(O)<sub>n</sub>NR<sub>B</sub>-, -OC(=S)S-, -NHC(=O)-, -NR<sub>B</sub>C(=O)-, -NHS(O)<sub>n</sub>-, -OSiH<sub>n</sub>( $C_{1-4}$ alkyl<sub>2-n</sub>)-, and -NR<sub>B</sub>S(O)<sub>n</sub>-;

Y and Z are independently selected at each occurrence from the group consisting of: 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, amino,  $C_{1-4}$ alkyl, -O( $C_{1-4}$ alkyl), -NH( $C_{1-4}$ alkyl), -N( $C_{1-4}$ alkyl)( $C_{1-4}$ alkyl), and -S(O)<sub>n</sub>(alkyl); and

n is 0, 1, or 2.

Particular embodiments of the invention include compounds and salts of Formula A in which

R<sub>X</sub> and R<sub>Y</sub> are the same or different and are independently selected from:

a) -(C=O)alkyl<sub>A</sub>, wherein alkyl<sub>A</sub> is a straight or branched alkyl group having from 1 to 8 carbon atoms;

b) straight, branched, or cyclic alkyl groups, including cycloalkyl(alkyl)groups, said alkyl groups having from 1 to 8 carbon atoms and optionally containing one or more double or triple bonds, each of which alkyl groups may be further substituted with one or more substituent(s) independently selected from:

i) hydroxy, halogen, amino, cyano, -O( $C_{1-4}$ alkyl), -NH( $C_{1-4}$ alkyl), and -NH( $C_{1-4}$ alkyl)( $C_{1-4}$ alkyl), and

ii) 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents independently selected from halogen, halo( $C_{1-4}$ alkyl, halo( $C_{1-4}$ alkoxy, oxo, hydroxy, amino,  $C_{1-4}$ alkyl, -O( $C_{1-4}$ alkyl), -NH( $C_{1-4}$ alkyl), -N( $C_{1-4}$ alkyl)( $C_{1-4}$ alkyl), and -S(O)<sub>n</sub>(alkyl), wherein said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) independently selected from N, O, and S, with the point of attachment being either carbon or nitrogen,

R<sub>1</sub> and R<sub>3</sub> are independently selected from C<sub>1-6</sub> alkyl<sub>1</sub>, (C<sub>3-7</sub>cycloalkyl<sub>1</sub>)C<sub>1-4</sub>alkyl<sub>1</sub>, -O(C<sub>3-7</sub>cycloalkyl<sub>1</sub>)C<sub>1-4</sub>alkyl<sub>1</sub>, halo(C<sub>1-6</sub>)alkyl<sub>1</sub>, -O(halo(C<sub>1-6</sub>)alkyl<sub>1</sub>), and -O(C<sub>1-6</sub>alkyl<sub>1</sub>),

where each said alkyl<sub>1</sub> is straight, branched, or cyclic and may contain 1 or more double or triple bonds, and is optionally substituted by one or more substituents independently chosen from halogen, hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino, and

where said C<sub>3-7</sub>cycloalkyl<sub>1</sub> is optionally substituted by one or more substituents independently chosen from halogen, hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino

Ar is phenyl, which is mono-, di-, or tri-substituted with R<sub>C</sub>;

R<sub>A</sub> and R<sub>B</sub>, which may be the same or different, are independently selected at each occurrence from the group consisting of:

hydrogen and straight, branched, or cyclic alkyl groups, including (cycloalkyl)alkyl groups, consisting of 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from oxo, hydroxy, halogen, nitro, cyano, C<sub>1-6</sub>alkoxy, -NH(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -NHC(=O)(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)C(=O)(C<sub>1-6</sub>alkyl), and Z;

R<sub>C</sub> is independently selected at each occurrence from halogen, cyano, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, hydroxy, amino, and C<sub>1-6</sub>alkyl substituted with 0-2 R<sub>D</sub>, C<sub>2-6</sub> alkenyl substituted with 0-2 R<sub>D</sub>, C<sub>2-6</sub>alkynyl substituted with 0-2 R<sub>D</sub>, C<sub>3-7</sub>cycloalkyl substituted with 0-2 R<sub>D</sub>, (C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl substituted with 0-2 R<sub>D</sub>, C<sub>1-6</sub>alkoxy substituted with 0-2 R<sub>D</sub>, -NH(C<sub>1-6</sub>alkyl) substituted with 0-2 R<sub>D</sub>, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl) each C<sub>1-4</sub>alkyl independently substituted with 0-2 R<sub>D</sub>, -XR<sub>A</sub>, and Y, with the proviso that at least one of the positions ortho or para to the point of attachment of Ar to the pyrimidine ring shown in Formula A is substituted;

R<sub>D</sub> is independently selected at each occurrence the group consisting of halogen, hydroxy, cyano, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, CO(C<sub>1-4</sub>alkyl), CONH(C<sub>1-4</sub>alkyl), CON(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), -XR<sub>A</sub>, and Y;

X is independently selected at each occurrence from the group consisting of -CH<sub>2</sub>-, -CHR<sub>B</sub>-, -O-, -C(=O)-, -C(=O)O-, -NH-, -NR<sub>B</sub>-, -C(=O)NH-, -C(=O)NR<sub>B</sub>-, -NHC(=O)-, and -NR<sub>B</sub>C(=O)-;

Y and Z are independently selected at each occurrence from the group consisting of: 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl); and

n is 0, 1, or 2.

Preferred compounds and salts of Formula A are those in which Ar is phenyl mono-, di-, or tri-substituted with R<sub>C</sub>, and R<sub>1</sub> and R<sub>3</sub> are independently selected from groups

(1) hydrogen, halogen, C<sub>1-4</sub>alkoxy, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, and

(2) C<sub>1-6</sub>alkyl and (C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl, wherein each member of group (2) is

unsubstituted or substituted by one to three substituents independently selected

from hydroxy, oxo, cyano, C<sub>1-4</sub>alkoxy, amino, and mono- or di(C<sub>1-4</sub>)alkylamino.


This class of embodiments of the inventions particularly includes compounds and salts in which

R<sub>X</sub> and R<sub>Y</sub>, which may be the same or different, are independently

selected at each occurrence from straight, branched, or cyclic alkyl groups, including

(cycloalkyl)alkyl groups, consisting of 1 to 8 carbon atoms, which may contain one or more double or triple bonds.

Other embodiments of the invention include compounds and salts of Formula A in which Ar is a phenyl group of the formula:




wherein L indicates a bond to the pyrimidine ring of Formula A

and the phenyl group is substituted at one, two, or three of positions 2, 4, and 6 positions of the phenyl ring with substituents independently selected from:

i) halogen, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, hydroxy, amino, C<sub>1-6</sub> alkyl, C<sub>1-6</sub>alkoxy, (C<sub>1-4</sub>alkoxy)C<sub>1-4</sub>alkoxy, and mono- or di(C<sub>1-4</sub>alkyl)amino,

ii) C<sub>1-6</sub> alkyl and C<sub>1-6</sub>alkoxy which are further substituted with a 3- to 7-membered carbocyclic and heterocyclic group, which is saturated, unsaturated, or aromatic, which 3- to 7-membered carbocyclic and heterocyclic group may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl).

Also provided by the invention are compounds and salts of Formula Ic




**Formula Ic**

wherein: R<sub>x</sub> is chosen from straight, branched, or cyclic alkyl groups, including cycloalkyl(alkyl) groups, having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from:

(a) hydroxy, halogen, amino, cyano, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -NH(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and

(b) 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, halo(C<sub>1-4</sub>)alkyl, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), wherein said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) selected from N, O, and S, with the point of attachment being either carbon or nitrogen. The remaining variables Ar, R<sub>1</sub> and R<sub>3</sub> are as defined for Formula I, for Formula Ia (preferred) or Formula A. The inventions particularly includes compounds and salts of Formula Ic in which Ar is as defined for Formula Ia and R<sub>1</sub> and R<sub>3</sub> are independently selected from the group consisting of hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, and halo(C<sub>1-4</sub>)alkyl.

Another embodiment of the invention includes compounds and salts of Formula B




**Formula B**

In which Ar is phenyl mono-, di-, or tri-substituted with R<sub>C</sub> (where R<sub>C</sub> carries the definition given in Formula Ia); R is selected from straight, branched, or cyclic alkyl groups, including (cycloalkyl)alkyl groups, which may contain 1 or more double or triple bonds, and which are optionally substituted by one or more substituents independently chosen from oxo, hydroxy, halogen, cyano, -O(C<sub>1-4</sub> alkyl), amino, -NH(C<sub>1-4</sub> alkyl), and -N(C<sub>1-4</sub> alkyl)(C<sub>1-4</sub> alkyl); R<sub>1</sub> is selected from hydrogen, halogen, cyano, C<sub>1-4</sub> alkyl, (C<sub>3-7</sub>cycloalkyl)C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, and -O(C<sub>1-4</sub>alkyl); and

R<sub>X</sub> and R<sub>Y</sub> are the same or different and are independently selected from:

- a) hydrogen,
- b) -(C=O)alkyl<sub>A</sub>, wherein alkyl<sub>A</sub> is a straight or branched alkyl group having from 1 to 8 carbon atoms;
- c) straight, branched, or cyclic alkyl groups, including cycloalkyl(alkyl) groups, having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from:
  - i) hydroxy, halogen, amino, cyano, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -NH(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and
  - ii) 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and -S(O)<sub>n</sub>(alkyl), wherein said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) independently selected from N, O, and S, with the point of attachment being either carbon or nitrogen.

Preferred compound of Formula B are those in which Ar is a phenyl group of the formula:




wherein L indicates a bond to the pyrimidine ring in Formula B

- 5 and the Ar phenyl group is substituted at one, two, or three of positions 2, 4, and 6 with substituents independently selected from (i) halogen, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, hydroxy, amino, C<sub>1-6</sub> alkyl, C<sub>1-6</sub>alkoxy, (C<sub>1-4</sub>alkoxy)C<sub>1-4</sub>alkoxy, and mono- or di(C<sub>1-4</sub>alkyl)amino, and (ii) C<sub>1-6</sub> alkyl and C<sub>1-6</sub>alkoxy which are further substituted with a 3- to 7-membered carbocyclic and heterocyclic group, which is saturated, unsaturated, or aromatic, which 3- to 7-membered carbocyclic and heterocyclic group may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl).

- Preferred compound and salts of this embodiment of the invention include compounds in which R<sub>x</sub> and R<sub>y</sub> are the same or different and are independently selected from the groups (a) hydrogen (with the proviso that R<sub>x</sub> and R<sub>y</sub> are not both hydrogen), (b) -(C=O)alkyl<sub>A</sub>, wherein alkyl<sub>A</sub> is a straight or branched alkyl group having from 1 to 8 carbon atoms; and (c) straight, branched, or cyclic alkyl groups, including cycloalkyl(alkyl), said straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, and containing zero, one or more double or triple bonds, each of which 1 to 8 carbon atoms may be further substituted with one or more substituent(s) independently selected from hydroxy, halogen, amino, cyano, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -NH(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl).

Additional preferred compounds of Formula B are those wherein Ar is a phenyl group of the formula:



wherein L indicates a bond to the pyrimidine ring in Formula B



and the Ar phenyl group is substituted at one, two, or three of positions 2, 4, and 6 with substituents independently selected from:

i) halogen, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, hydroxy, amino, C<sub>1-6</sub> alkyl, C<sub>1-6</sub>alkoxy, (C<sub>1-4</sub>alkoxy)C<sub>1-4</sub>alkoxy, and mono- or di(C<sub>1-4</sub>alkyl)amino,

ii) C<sub>1-6</sub> alkyl and C<sub>1-6</sub>alkoxy which are further substituted with a 3- to 7-membered carbocyclic and heterocyclic group, which is saturated, unsaturated, or aromatic, which 3- to 7-membered carbocyclic and heterocyclic group may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl);


R<sub>X</sub> and R<sub>Y</sub> are the same or different and are independently selected from the group consisting of:

a) hydrogen (with the proviso that R<sub>X</sub> and R<sub>Y</sub> are not both hydrogen),

b) -(C=O)alkyl<sub>A</sub>, wherein alkyl<sub>A</sub> is a straight or branched alkyl group having from 1 to 8 carbon atoms;

c) straight, branched, or cyclic alkyl groups, including cycloalkyl(alkyl) groups, said straight, branched, or cyclic alkyl groups have from 1 to 8 carbon atoms and may contain one or more double or triple bonds.

The invention further provides compounds and salts of Formula C



**Formula C**

wherein R, R<sub>1</sub>, R<sub>X</sub>, and R<sub>Y</sub> carry the definitions set forth for Formula B and

q is an integer from 1 to 4;

G is hydrogen, hydroxy, C<sub>1-6</sub>alkoxy, -NH(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), or a 3- to 7-

membered carbocyclic or heterocyclic group which is saturated, unsaturated, or aromatic,


which is unsubstituted or substituted with one or more substituents independently

selected from halogen, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, oxo, hydroxy, amino, C<sub>1-4</sub>alkyl,

-O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and -S(O)<sub>n</sub>(alkyl), wherein said 3- to 7-membered heterocyclic group contains one or more heteroatom(s) independently selected from N, O, and S, with the point of attachment being either carbon or nitrogen;

- 5 J and K are independently selected from halogen, cyano, halo(C<sub>1-4</sub>alkyl), halo(C<sub>1-4</sub>alkoxy), hydroxy, amino, C<sub>1-6</sub> alkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, (C<sub>1-4</sub>alkoxy) C<sub>1-4</sub>alkoxy, and mono- or di(C<sub>1-4</sub>alkyl)amino.

In yet another embodiment the invention provides compounds and salts of Formula D



**Formula D**

wherein R and R<sub>1</sub> carry the definitions set forth for Formula B:

Q is hydrogen, C<sub>3-7</sub> cycloalkyl, pyrrolidinyl, piperidinyl, morpholino, or piperazinyl;


- 15 q is an integer from 1 to 4;

G is hydrogen, hydroxy, C<sub>1-6</sub>alkoxy, -NH(C<sub>1-6</sub>alkyl), -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), or a 3- to 7-membered carbocyclic or heterocyclic group, which is saturated, unsaturated, or aromatic, which is unsubstituted or substituted with one or more substituents independently selected from halogen, halo(C<sub>1-4</sub>alkyl), halo(C<sub>1-4</sub>alkoxy), oxo, hydroxy, amino, C<sub>1-4</sub>alkyl, -O(C<sub>1-4</sub>alkyl), -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)(C<sub>1-4</sub>alkyl), and -S(O)<sub>n</sub>(alkyl), wherein said 3- to 7-membered heterocyclic group contains one or more heteroatom(s) independently selected from N, O, and S, with the point of attachment being either carbon or nitrogen;

J and K are independently selected from halogen, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, hydroxy, amino, C<sub>1-6</sub> alkyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, (C<sub>1-4</sub>alkoxy) C<sub>1-4</sub>alkoxy, and mono- or di(C<sub>1-4</sub>alkyl)amino; and

R<sub>X</sub> and R<sub>Y</sub> are the same or different and are independently selected from hydrogen (with the proviso that R<sub>X</sub> and R<sub>Y</sub> are not both hydrogen) and straight, branched, or cyclic alkyl groups having from 1 to 6 carbon atoms, which alkyl groups may contain one or more double or triple bonds.

Other compound and salts of the invention are those having Formula E



**Formula E**

wherein R<sub>1</sub>, R<sub>3</sub>, and Ar may carry the definitions set forth for Formula Ia and A is NH, N(C<sub>1-6</sub>-alkyl), CH<sub>2</sub>, CH(C<sub>1-6</sub>-alkyl) or O.

Compounds of the invention are useful in treating a variety of conditions including affective disorders, anxiety disorders, stress disorders, eating disorders, and drug addiction.

Affective disorders include all types of depression, bipolar disorder, cyclothymia, and dysthymia.

Anxiety disorders include generalized anxiety disorder, panic, phobias and obsessive-compulsive disorder.

Stress-related disorders include post-traumatic stress disorder, hemorrhagic stress, stress-induced psychotic episodes, psychosocial dwarfism, stress headaches, stress-induced immune systems disorders such as stress-induced fever, and stress-related sleep disorders.

Eating disorders include anorexia nervosa, bulimia nervosa, and obesity.

Modulators of the CRF receptors may also be useful in the treatment of a variety of neurological disorders including supranuclear palsy, AIDS related dementias, multiinfarct dementia, neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, and Huntington's disease, head trauma, spinal cord trauma, ischemic neuronal damage, amyotrophic lateral sclerosis, disorders of pain perception such as fibromyalgia and epilepsy.

Additionally compounds of Formula I are useful as modulators of the CRF receptor in the treatment of a number of gastrointestinal, cardiovascular, hormonal, autoimmune and inflammatory conditions. Such conditions include irritable bowel syndrome, ulcers, Crohn's disease, spastic colon, diarrhea, post operative ilius and colonic hypersensitivity associated with psychopathological disturbances or stress, hypertension, tachycardia, congestive heart failure, infertility, euthyroid sick syndrome, inflammatory conditions effected by rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies.

Compounds of Formula I are also useful as modulators of the CRF<sub>1</sub> receptor in the treatment of animal disorders associated with aberrant CRF levels. These conditions include porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs, psychosocial dwarfism and hypoglycemia.

Typical subjects to which compounds of the invention may be administered will be mammals, particularly primates, especially humans. For veterinary applications, a wide variety of subjects will be suitable, e.g. livestock such as cattle, sheep, goats, cows, swine and the like; poultry such as chickens, ducks, geese, turkeys, and the like; and domesticated animals particularly pets such as dogs and cats. For diagnostic or research applications, a wide variety of mammals will be suitable subjects including rodents (e.g. mice, rats, hamsters), rabbits, primates, and swine such as inbred pigs and the like. Additionally, for *in vitro* applications, such as *in vitro* diagnostic and research applications, body fluids and cell samples of the above subjects will be suitable for use such as mammalian, particularly primate such as human, blood, urine or tissue samples, or blood urine or tissue samples of the animals mentioned for veterinary applications.

The CRF binding compounds provided by this invention and labeled derivatives thereof are also useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the CRF receptor.

Labeled derivatives the CRF antagonist compounds provided by this invention are also useful as radiotracers for positron emission tomography (PET) imaging or for single photon emission computerized tomography (SPECT).

The present invention also pertains to methods of inhibiting the binding of CRF to CRF receptors which methods involve contacting a solution containing compound of the invention with cells expressing CRF receptors, wherein the compound is present in the solution at a

concentration sufficient to inhibit CRF binding to CRF receptors *in vitro*. This method includes inhibiting the binding of CRF to CRF receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to inhibit the binding of CRF to CRF receptors *in vitro*. In one embodiment, such methods are useful in treating physiological disorders associated with excess concentrations of CRF. The amount of a compound that would be sufficient to inhibit the binding of a CRF to the CRF receptor may be readily determined via a CRF receptor binding assay (see Example 96), or from the EC<sub>50</sub> of a CRF receptor functional assay, such as a standard assay of CRF receptor mediated chemotaxis. The CRF receptors used to determine *in vitro* binding may be obtained from a variety of sources, for example from cells that naturally express CRF receptors, e.g. IMR32 cells or from cells expressing cloned human CRF receptors.

The present invention also pertains to methods for altering the activity of CRF receptors, said method comprising exposing cells expressing such receptors to an effective amount of a compound of the invention, wherein the compound is present in the solution at a concentration sufficient to specifically alter the signal transduction activity in response to CRF in cells expressing high levels of CRF1 receptors *in vitro*. This method includes altering the signal transduction activity of CRF receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to alter the signal transduction activity in response to CRF in cells expressing high levels of CRF1 *in vitro*. The amount of a compound that would be sufficient to alter the signal transduction activity in response to CRF receptors may be determined via an assay of CRF receptor mediated signal transduction, such as an assay wherein the binding of CRF to a cell surface CRF receptor effects a changes in reporter gene expression.

The present invention also pertains to packaged pharmaceutical compositions for treating disorders responsive to C5a receptor modulation, e.g., eating disorders, depression or stress. The packaged pharmaceutical compositions include a container holding a therapeutically effective amount of at least one CRF1 receptor modulator as described supra and instructions for using the treating disorder responsive to CRF1 receptor modulation in the patient.

### **Definitions**

The compounds herein described may have one or more asymmetric centers or planes. Compounds of the present invention containing an asymmetrically substituted atom may be

isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms (racemates), by asymmetric synthesis, or by synthesis from optically active starting materials. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography, using, for example a chiral HPLC column. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral (enantiomeric and diastereomeric), and racemic forms, as well as all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

When any variable occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R<sup>\*</sup>, then said group may optionally be substituted with up to two R<sup>\*</sup> groups and R<sup>\*</sup> at each occurrence is selected independently from the definition of R<sup>\*</sup>. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

Compounds of Formula I include, but are not limited to, compounds of Formula Ia, Ic, A, B, C, D, and E.

As indicated above, various substituents of the various formulae are "optionally substituted", including Ar, R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> of Formula I and subformulae thereof, and such substituents as recited in the sub-formulae such as Formula I and subformulae, e.g. Formula Ia, Ic, A, B, C, D, and E and the like. The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valence is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers, By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include <sup>11</sup>C, <sup>13</sup>C, and <sup>14</sup>C.

When substituted, those substituents (Ar, R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub>) may be substituted by other than hydrogen at one or more available positions, typically 1 to 3 or 4 positions, by one or more suitable groups such as those disclosed herein. Suitable groups that may be present on a "substituted" Ar, R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> group or other substituent include e.g. halogen such as fluoro, chloro, bromo and iodo; cyano; hydroxyl; nitro; azido; alkanoyl such as a C<sub>1-6</sub> alkanoyl group such as acyl and the like; carboxamido; alkyl groups, including cycloalkyl groups, having 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5, or 6 carbon atoms; alkenyl and alkynyl groups including groups having one or more unsaturated linkages and from 2 to about 12 carbon, or 2, 3, 4, 5 or 6 carbon atoms; alkoxy groups having those having one or more oxygen linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5 or 6 carbon atoms; aryloxy such as phenoxy; alkylthio groups including those moieties having one or more thioether linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5 or 6 carbon atoms; alkylsulfinyl groups including those moieties having one or more sulfinyl linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5, or 6 carbon atoms; alkylsulfonyl groups including those moieties having one or more sulfonyl linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5, or 6 carbon atoms; aminoalkyl groups such as groups having one or more N atoms and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5 or 6 carbon atoms; carbocyclic aryl having 6 or more carbons, particularly phenyl (e.g. an Ar group being a substituted or unsubstituted biphenyl moiety); aralkyl having 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms, with benzyl being a preferred group; aralkoxy having 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms, with O-benzyl being a preferred group; or a saturated, unsaturated, or aromatic heterocyclic group having 1 to 3 separate or fused rings with 3 to about 8 members per ring and one or more N, O or S atoms, e.g. coumarinyl, quinolinyl, pyridyl, pyrazinyl, pyrimidyl, furyl, pyrrolyl, thienyl, thiazolyl, oxazolyl, imidazolyl, indolyl, benzofuranyl, benzothiazolyl, tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, morpholino and pyrrolidinyl. Such heterocyclic groups may be further substituted, e.g. with hydroxy, alkyl, halogen and amino.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups, having the specified number of carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl. Preferred alkyl groups are C<sub>1</sub>-C<sub>6</sub> alkyl groups. Especially preferred alkyl groups are methyl, ethyl, propyl, butyl, 3-pentyl. The term C<sub>1-4</sub> alkyl as used herein includes

alkyl groups consisting of 1 to 4 carbon atoms, which may contain a cyclopropyl moiety. Suitable examples are methyl, ethyl, and cyclopropylmethyl.

"Cycloalkyl" is intended to include saturated ring groups, having the specified number of carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. Cycloalkyl groups typically will have 3 to about 8 ring members.

In the term "(C<sub>3-6</sub> cycloalkyl)C<sub>1-4</sub> alkyl", as defined above, the point of attachment is on the alkyl group. This term encompasses, but is not limited to, cyclopropylmethyl, cyclohexylmethyl, cyclohexylmethyl.

"Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration comprising one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl and propenyl. Alkenyl groups typically will have 2 to about 12 carbon atoms, more typically 2 to about 8 carbon atoms.

"Alkynyl" is intended to include hydrocarbon chains of either a straight or branched configuration comprising one or more triple carbon-carbon bonds which may occur in any stable point along the chain, such as ethynyl and propynyl. Alkynyl groups typically will have 2 to about 12 carbon atoms, more typically 2 to about 8 carbon atoms.

"Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example -C<sub>v</sub>F<sub>w</sub> where v = 1 to 3 and w = 1 to (2v+1)). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl. Typical haloalkyl groups will have 1 to about 8 carbon atoms, more typically 1 to about 6 carbon atoms.

"Alkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, 2-butoxy, t-butoxy, n-pentoxy, 2-pentoxy, 3-pentoxy, isopentoxy, neopentoxy, n-hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy. Alkoxy groups typically have 1 to about 12 carbon atoms, more typically 1 to about 8 carbon atoms.

As used herein, the term "alkylthio" includes those groups having one or more thioether linkages and suitably from 1 to about 12 carbon atoms, more typically 1 to about 8 carbon atoms, still more typically 1 to about 6 carbon atoms.



As used herein, the term "alkylsulfinyl" includes those groups having one or more sulfoxide (SO) linkage groups and suitably from 1 to about 12 carbon atoms, more typically 1 to about 8 carbon atoms, still more typically 1 to about 6 carbon atoms.

As used herein, the term "alkylsulfonyl" includes those groups having one or more sulfonyl (SO<sub>2</sub>) linkage groups and suitably from 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms, still more typically 1 to about 6 or 8 carbon atoms.

As used herein, the term "alkylamino" includes those groups having one or more primary, secondary and/or tertiary amine groups and suitably from 1 to about 12 carbon atoms, more typically 1 to about 8 carbon atoms, still more typically 1 to about 6 carbon atoms.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo; and "counter-ion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

As used herein, "carbocyclic group" is intended to mean any stable 3- to 7-membered monocyclic or bicyclic or 7-to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl.

As used herein, the term "heterocyclic group" is intended to mean a stable 5-to 7-membered monocyclic or bicyclic or 7-to 10-membered bicyclic heterocyclic ring which is saturated, partially unsaturated, or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The term or "heterocycloalkyl" is used to refer to saturated heterocyclic groups.

The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used

herein, the term "aromatic heterocyclic system" is intended to mean a stable 5-to 7-membered monocyclic or bicyclic or 7-to 10membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolinyl, carbazolyl, NH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2*H*,6*H*-1,5,2-dithiazinyl, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolinyl, imidazolyl, 1*H*-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, 3*H*-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl;- 1,2,5oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2*H*-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4*H*-quinoliziny, quinoxaliny, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6*H*-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

Preferred heterocyclic groups include, but are not limited to, pyridinyl, pyrimidinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrrolidinyl, morpholinyl, piperidinyl, piperazinyl, and imidazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

As used herein, the term "carbocyclic aryl" includes groups that contain 1 to 3 separate or fused rings and from 6 to about 18 ring atoms, without hetero atoms as ring members.

Specifically preferred carbocyclic aryl groups include phenyl, and naphthyl including 1-naphthyl and 2-naphthyl.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings or animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic,  $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$  where  $n$  is 0-4, and the like. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, p. 1418 (1985).

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to Formula I *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of formula I are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of formula I wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of formula I is administered to a mammalian subject, cleaves to form a

free hydroxyl, free amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula I, and the like.

Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. A stable compound or stable structure is meant to imply a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an effective therapeutic agent. The term "therapeutically effective amount" of a compound of this invention means an amount effective to antagonize abnormal level of CRF or treat the symptoms of affective disorder, anxiety or depression in a host.

### **Pharmaceutical Preparations**

The compounds of general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrathecal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for

example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

5 Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable  
10 for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of  
15 ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example  
20 ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard  
25 paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting  
30 agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents

and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compounds of general Formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

Frequency of dosage may also vary depending on the compound used and the particular disease treated. However, for treatment of most CNS disorders, a dosage regimen of 4 times daily or less is preferred. For the treatment of stress and depression a dosage regimen of 1 or 2 times daily is particularly preferred.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

Preferred compounds of the invention will have certain pharmacological properties. Such properties include, but are not limited to oral bioavailability, low toxicity, low serum protein binding and desirable *in vitro* and *in vivo* half-lives. Penetration of the blood brain barrier for compounds used to treat CNS disorders is necessary, while low brain levels of compounds used to treat peripheral disorders are often preferred.

Assays may be used to predict these desirable pharmacological properties. Assays used to predict bioavailability include transport across human intestinal cell monolayers, including Caco-2 cell monolayers. Toxicity to cultured hepatocytes may be used to predict compound toxicity. Penetration of the blood brain barrier of a compound in humans may be predicted from the brain levels of the compound in laboratory animals given the compound intravenously.

Serum protein binding may be predicted from albumin binding assays. Such assays are described in a review by Oravcová, et al. (Journal of Chromatography B (1996) volume 677, pages 1-27).

Compound half-life is inversely proportional to the frequency of dosage of a compound. *In vitro* half-lives of compounds may be predicted from assays of microsomal half-life as described by Kuhnz and Gieschen (Drug Metabolism and Disposition, (1998) volume 26, pages 1120-1127).

As discussed above, preferred arylpyrimidines of the invention exhibit good activity in standard *in vitro* CRF receptor binding assays, specifically the assay as specified in Example 96, which follows. References herein to “standard *in vitro* receptor binding assay” are intended to refer to that protocol as defined in Example 96 which follows. Generally preferred compounds preferred arylpyrimidines of the invention have an IC<sub>50</sub> of about 1 micromolar or less, still more preferably and IC<sub>50</sub> of about 100 nanomolar or less even more preferably an IC<sub>50</sub> of about 10 nanomolar or less or even 1 nanomolar or less in such a defined standard *in vitro* CRF receptor binding assay as exemplified by Example 96 which follows.

## EXAMPLES


### Preparation of Arylpyrimidines

The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include but are not limited to those methods described below. Each of the references cited below are hereby incorporated herein by reference. Preferred methods for the preparation of compounds of the present invention include, but are not limited to, those described in Scheme I to Scheme IV. Those who are skilled in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention. All references cited herein are hereby incorporated in their entirety herein by reference. The following abbreviations are used herein:

|                   |                                 |
|-------------------|---------------------------------|
| AcOH              | acetic acid                     |
| DMF               | N,N-dimethylformamide           |
| Et <sub>2</sub> O | diethyl ether                   |
| EtOAc             | ethyl acetate                   |
| EtOH              | ethanol                         |
| NaH               | sodium hydride                  |
| NaHMDS            | sodium bis(trimethylsilyl)amide |
| THF               | tetrahydrofuran                 |
| EX#               | example number                  |




### Scheme I (Method A)



5 According to the general method A, wherein  $R_1$  and  $R_3$  are as defined for formula I and Hal represents a halogen atom, suitably chloride or bromide. Compounds of formula IV can be prepared according to a known literature procedure (Ref: *Journal of Organic Chemistry* **1983**, 48, 1060). Reduction of the nitro group in IV may be accomplished by a variety of methods known in the art, including hydrogenation with hydrogen and transition metal catalysts or the use of sodium hydrosulfite in aqueous solutions to give V. The amino pyrimidine V may be transformed into VI by reductive amination using aldehydes and reducing agents such as sodium triacetoxyborohydride in inert solvents. The halopyrimidine VI can be converted to arylpyrimidine II by a transition metal-catalyzed coupling reaction with a metalloaryl reagent (Ar-[M]). More commonly employed reagent/catalyst pairs include aryl boronic acid/palladium(0) (Suzuki reaction; N. Miyaura and A. Suzuki, *Chemical Review* 1995, 95, 2457), aryl trialkylstannane/palladium(0) (Stille reaction; T. N. Mitchell, *Synthesis* 1992, 803), arylzinc/palladium(0) and aryl Grignard/nickel(II). Palladium(0) represents a catalytic system made of a various combination of metal/ligand pair which includes, but not limited to, tetrakis(triphenylphosphine)palladium(0), palladium(II) acetate/tri(*o*-tolyl)phosphine, 15 tris(dibenzylideneacetone)dipalladium(0)/tri-*tert*-butylphosphine and dichloro[1,1'-bis(diphenylphosphine)ferrocene]palladium(0). Nickel(II) represents a nickel-containing catalyst such as [1,2-bis(diphenylphosphino)ethane]dichloronickel(II) and [1,3-bis(diphenylphosphino)propane]dichloronickel(II).


### Scheme II (Method B)



By changing the sequence of events in Scheme I, but using same the methods described in

5 Method A, compounds of the formula II can also be prepared as outlined in Scheme II

### Scheme III (Method C)




10 An alternative method for introducing the substituents  $R_A$  and  $R_B$  to give compounds of the formula II is outlined in Scheme III and may be accomplished by a variety of methods known in the art. These include reaction of the amine VIII with acid chlorides or anhydrides in the presence of bases such as but not limited to triethylamine or pyridine in inert solvents such as dichloromethane or toluene. The N-H group is then deprotonated by a strong base such as but not limited to alkali metal hydride, alkali metal amide, or alkali metal alkoxide in inert solvents such as but not limited to THF, DMF, or methyl sulfoxide.

15

Alkylation may be conducted using alkyl halide, suitably bromide or iodide, at temperatures ranging from  $0^\circ C$  to  $100^\circ C$ . Reduction of the amide IX with reducing agents such as but not limited to lithium aluminum hydride, borane or diisobutylaluminum hydride in inert solvents such as but not limited to THF, ether, or toluene furnishes compounds of the formula II.

20

#### Scheme IV (Method D)



Yet another method for preparing compounds of formula **II** is illustrated in the Scheme IV. Treatment of the amine **VIII** with base such as but not limited to alkali metal hydride, alkali metal amide, alkali metal alkoxide or alkali metal carbonates in inert solvents such as but not limited to THF, DMF, methyl sulfoxide or acetonitrile with or without the addition of alkali metal iodide, followed by alkylation with alkyl halide, suitably bromide or iodide, or sulfonate at temperatures ranging from  $0^\circ\text{C}$  to  $100^\circ\text{C}$  furnishes compounds of the formula **II**.

#### EXAMPLES

The preparation of the compounds of the present invention is illustrated further by the following examples, which are not to be construed as limiting the invention in scope or spirit to the specific procedures and compounds described in them.

Commercial reagents are used without further purification. room or ambient temperature refers to  $20$  to  $25^\circ\text{C}$ . Concentration *in vacuo* implies the use of a rotary evaporator. TLC refers to thin layer chromatography. Proton nuclear magnetic resonance ( $^1\text{H}$  NMR) spectral data are obtained at  $300$  or  $400$  MHz. Mass spectral data are obtained either by CI or APCI methods.

##### Example 1

**[2-(2,4-dimethoxyphenyl)-4-methoxy-6-methylpyrimidin-5-yl]dipropylamine** [Formula I:  $\text{Ar} = 2,4\text{-dimethoxyphenyl}$ ;  $\text{R}_1 = \text{OCH}_3$ ;  $\text{R}_2 = \text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$ ;  $\text{R}_3 = \text{CH}_3$ ]


A. To a solution of 2-chloro-4-methoxy-6-methyl-5-nitropyrimidine (1.01 g, 5.00 mmol) in THF (20 mL) and water (20 mL) is added sodium hydrosulfite (8.6 g, 50.0 mmol). The mixture is stirred at room temperature for 2.5 h, diluted with saturated aqueous sodium bicarbonate and extracted with ethyl acetate. Combined extracts are washed with brine, dried (sodium sulfate), filtered and concentrated to give 2-chloro-4-methoxypyrimidine-5-ylamine (600 mg):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  2.38 (s, 3H), 4.02 (s, 3H).

B. To a solution of 2-chloro-4-methoxypyrimidine-5-ylamine (400 mg, 2.3 mmol) in 1,2-dichloroethane (60 mL) is added propionaldehyde (700 mg, 12 mmol) and glacial acetic acid (660 mg). After 10 minutes sodium triacetoxymethylborohydride (2.5g, 12 mmol) is added in one  
5 portion. After 3 h the volatiles are removed by rotary evaporation. The residue is partitioned between ethyl acetate and saturated aqueous sodium bicarbonate, the layers are separated and the aqueous layer further extracted with ethyl acetate. The combined organics are washed with water, brine, dried (sodium sulfate), filtered and concentrated to give (2-chloro-4-methoxy-6-methylpyrimidine-5-yl)dipropylamine (566 mg): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 0.85 (t, 6H), 1.3  
10 (m, 4H), 2.4 (s, 3H), 2.82 (t, 4H), 4.0 (s, 3H); MS (CI) 258.

C. A stirred solution of (2-chloro-4-methoxy-6-methylpyrimidine-5-yl)dipropylamine (380 mg; 1.47 mmol) and tetrakis(triphenylphosphine)palladium(0) (33 mg; 2 mol%) in ethyleneglycol dimethyl ether (8 mL) is stirred at room temperature for 15 min, then 2,4-  
15 dimethoxybenzeneboronic acid (1.76 mmol) and an aqueous solution of sodium carbonate (1.0 M, 4 mL) are added sequentially. The mixture is heated to 75°C with stirring for 1.5 h, then diluted with 0.1 N sodium hydroxide and extracted twice with 1:1 hexane-ethyl ether. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed on silica (1:1 hexane-ether) to give the title compound (0.50 g): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 0.85  
20 (t, 6H), 1.4 (m, 4H), 2.58 (s, 3H), 2.9 (t, 4H), 3.82 (s, 3H), 3.84 (s, 3H), 4.0 (s, 3H), 6.58 (m, 2H), 7.8 (d, 1H); MS (CI) 360.

The **EX#s 2-6b** in the Table I may be prepared following the methods described in Example 1.

Table I



| EX# | Ar                          | <sup>1</sup> H NMR (CDCl <sub>3</sub> )   | MS (CI) | Name  |
|-----|-----------------------------|---|---------|---|
| 2   | 2-chlorophenyl              | 0.85 (t, 6H), 1.38 (m, 4H), 2.58 (s, 3H), 2.92 (t, 4H), 4.05 (s, 3H), 7.40 (m, 2H), 8.3 (d, 1H), 8.4 (br s, 1H)                         | 334     | [2-(2-chlorophenyl)-4-methoxy-6-methylpyrimidin-5-yl]dipropylamine              |
| 3   | 2,4-dichlorophenyl          | 0.82 (t, 6H), 1.4 (m, 4H), 2.58 (s, 3H), 2.9 (t, 4H), 4.0 (s, 3H), 7.24 (d, 1H), 7.50 (d, 1H), 7.6 (d, 1H)                              | 368     | [2-(2,4-dichlorophenyl)-4-methoxy-6-methylpyrimidin-5-yl]dipropylamine          |
| 4   | 4-chloro-2-methoxyphenyl    | 0.85 (t, 6H), 1.4 (m, 4H), 2.56 (t, 4H), 3.85 (s, 3H), 4.05 (s, 3H), 6.9 (d, 1H), 7.3 (d, 1H), 7.75 (d, 1H)                             | 364     | [2-(2-methoxy-4-chlorophenyl)-4-methoxy-6-methylpyrimidin-5-yl]dipropylamine    |
| 5   | 2-methoxy-4-isopropylphenyl | 0.85 (t, 6H), 1.2 (d, 6H), 1.4 (m, 4H), 3.90 (m, 5H), 3.82 (s, 3H), 4.00 (s, 3H), 6.95 (d, 1H), 7.22 (d, 1H), 7.55 (d, 1H)              | 372     | [2-(2-methoxy-4-isopropylphenyl)-4-methoxy-6-methylpyrimidin-5-yl]dipropylamine |
| 6   | 2,4-dimethoxyphenyl         | 0.85 (t, 6H), 1.4 (m, 4H), 2.58 (s, 3H), 2.92 (t, 4H), 3.82 (s, 3H), 3.85 (m 2H), 4.02 (s, 3H), 6.90 (d, 1H), 6.95 (d, 1H), 7.3 (d, 1H) | 360     | [2-(2,4-dimethoxyphenyl)-4-methoxy-6-methyl pyrimidin-5-yl]dipropylamine        |
| 6a  | 2,6-dimethoxyphenyl         |   |         | [4-Methoxy-2-(2,6-dimethoxyphenyl)-6-methyl-pyrimidin-5-yl]-dipropyl-amine      |

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

|    |                                    |  |  |  |
|----|------------------------------------|--|--|--|
| 6b | 2-methoxy-6-trifluoromethoxyphenyl |  |  | [4-Methoxy-2-(2-methoxy-6-trifluoromethoxy-phenyl)-6-methyl-pyrimidin-5-yl]-dipropyl-amine |
|----|------------------------------------|--|--|--|

### Example 7

#### [4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]dipropylamine

[Formula I: Ar=6-methoxy-2,4-dimethylphenyl; R<sub>1</sub>=OCH<sub>3</sub>; R<sub>2</sub>=N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>; R<sub>3</sub>=CH<sub>3</sub>]

A. A solution of 2-chloro-4-methoxy-6-methyl-5-nitropyrimidine (305 mg, 1.5 mmol) and  
5 tetrakis(triphenylphosphine)palladium(0) (33 mg; 2 mol%) in ethyleneglycol dimethyl ether (8 mL) is stirred at room temperature for 15 min, then 2,4-dimethyl-6-methoxybenzeneboronic acid (1.76 mmol) and an aqueous solution of sodium carbonate (1.0 M, 4 mL) are added sequentially. The mixture is heated to 75°C with stirring for 1.5 h, then diluted with 0.1 N sodium hydroxide and extracted twice with 1:1 hexane-ethyl ether. Combined extracts are dried (sodium sulfate),  
10 filtered, concentrated and chromatographed on silica (4:1 hexane-ether) to give 3-methoxy-2-(4-methoxy-6-methyl-5-nitropyrimidin-2-yl)-1,5-dimethylbenzene (0.36 g): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) 2.08 (s, 3H), 2.35 (s, 3H), 2.58 (s, 3H), 3.7 (s, 3H), 4.05 (s, 3H), 6.65 (s, 1H), 6.72; MS (CI) 304.

15 B. To a solution of 3-methoxy-2-(4-methoxy-6-methyl-5-nitropyrimidin-2-yl)-1,5-dimethylbenzene (1.51 g, 5.00 mmol) in THF (20 mL) and water (20 mL) is added sodium hydrosulfite (8.6 g, 50.0 mmol). The mixture is stirred at room temperature for 14 h, diluted with saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. Combined extracts are washed with brine, dried (sodium sulfate), filtered and concentrated to give 4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidine-5-ylamine (1.05 g): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) 2.05 (s, 3H), 2.34 (s, 3H), 2.38 (s, 3H), 3.60 (br s, 2H), 3.68 (s, 3H), 4.00 (s, 3H), 6.60 (s, 1H), 6.64 (s, 1H).  
20

C. To a solution of 4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidine-5-ylamine (50 mg, 0.2 mmol) in 1,2-dichloroethane (5 mL) is added propionaldehyde (70 mg, 1.2 mmol) and glacial acetic acid (60 mg). After 10 minutes sodium triacetoxyborohydride (250 mg, 1.2 mmol) is added in one portion. The solution is stirred at room temperature for 1 h and the volatiles removed by rotary evaporation. The residue is partitioned between ethyl acetate and saturated aqueous sodium bicarbonate, the layers are separated and the aqueous layer further  
25 extracted with ethyl acetate. The combined organics are washed with water, brine, dried (sodium sulfate), filtered, concentrated and chromatographed on silica (4:1 hexane-ether) to give to give  
30


the title compound (56 mg):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  0.85 (t, 6H), 1.4 (m, 4H), 2.02 (s, 3H), 2.34 (s, 3H), 2.56 (s, 3H), 2.92 (t, 4H), 3.84 (s, 3H), 3.94 (s, 3H), 6.62 (s, 1H), 6.65 (s, 1H); MS (CI) 358.

The EX#s 8-20 in the Table II may be prepared following the methods described in Example 7.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200





Table II



| EX# | Ar                           | R <sub>1</sub> | R <sub>2</sub> | <sup>1</sup> H NMR (CDCl <sub>3</sub> )  | MS (CI) | NAME   |
|-----|------------------------------|----------------|----------------|--|---------|--|
| 8   | 2-methoxy-4,6-dimethylphenyl | Et             | Me             | 0.85 (t, 6H), 1.22 (t, 3H), 1.38 (m, 4H), 2.05 (s, 3H), 2.38 (s, 3H), 2.85 (m, 6H), 3.75 (s, 3H), 3.84 (s, 3H), 6.62 (s, 1H), 6.68 (s, 1H) | 372     | [2-(2-methoxy-4,6-dimethylphenyl)-4-methoxy-6-ethylpyrimidin-5-yl] dipropylamine |
| 9   | 2,4,6-trimethylphenyl        | Me             | Me             | 0.80 (t, 6H), 1.38 (m, 4H), 2.12 (s, 6H), 2.30 (s, 3H), 2.55 (s, 3H), 2.85 (m, 4H), 3.95 (s, 3H), 6.90 (s, 2H)                             | 342     | [2-(2,4,6-trimethylphenyl)-4-methoxy-6-methylpyrimidin-5-yl] dipropylamine       |
| 10  | 2,4,6-trimethylphenyl        | Et             | Me             | 0.85 (t, 6H), 1.22 (t, 3H), 1.40 (m, 4H), 2.15 (s, 6H), 2.30 (s, 3H), 2.90 (m, 6H), 3.92 (s, 3H), 6.9 (s, 2H)                              | 356     | [2-(2,4,6-trimethylphenyl)-4-methoxy-6-ethylpyrimidin-5-yl] dipropylamine        |


|     |                                    |                   |                                   |   |     |   |
|-----|------------------------------------|-------------------|-----------------------------------|---|-----|---|
| 11  | 2-methoxy-4,6-dimethylphenyl       | Me                | Et                                | 0.89 (t, 6H), 1.35 (t, 3H), 1.37 (m, 4H), 2.05 (s, 3H), 2.32 (s, 3H), 2.52 (s, 3H), 2.94 (t, 4H), 3.73 (s, 3H), 4.40 (q, 2H), 6.62 (s, 1H), 6.66 (s, 1H)    | 372 | [2-(2-methoxy-4,6-dimethylphenyl)-4-ethoxy-6-methyl pyrimidin-5-yl] dipropylamine           |
| 11a | 2-methoxy-6-trifluoromethoxyphenyl | Me                | Et                                |   |     | [2-(2-methoxy-6-trifluoromethoxyphenyl)-4-ethoxy-6-methyl pyrimidin-5-yl] dipropylamine     |
| 12  | 2-methoxy-4,6-dimethylphenyl       | Me                | CH <sub>2</sub> CH <sub>2</sub> F | 0.89 (t, 6H), 1.40 (m, 4H), 2.04 (s, 3H), 2.33 (s, 3H), 2.55 (s, 3H), 2.96 (t, 4H), 3.72 (s, 3H), 4.64 (m, 2H), 4.66 (m, 2H), 6.62 (s, 1H), 6.67 (s, 1H)    | 390 | [2-(2-methoxy-4,6-dimethylphenyl)-4-(2-fluoroethoxy)-6-methyl pyrimidin-5-yl] dipropylamine |
| 13  | 2-methoxy-4,6-dimethylphenyl       | Me                | CH(CH <sub>3</sub> ) <sub>2</sub> | 0.89 (t, 6H), 1.31 (d, 6H), 1.36 (m, 4H), 2.06 (s, 3H), 2.33 (s, 3H), 2.51 (s, 3H), 2.94 (t, 4H), 3.73 (s, 3H), 5.38 (hept, 1H), 6.62 (s, 1H), 6.67 (s, 1H) | 344 | [2-(2-methoxy-4,6-dimethylphenyl)-4-isopropoxy-6-methyl pyrimidin-5-yl] dipropylamine       |
| 14  | 2-methoxy-4,6-dimethylphenyl       | CH <sub>2</sub> F | Me                                | 0.88 (t, 6H), 1.38 (m, 4H), 2.07 (s, 3H), 2.34 (s, 3H), 2.94 (t, 4H), 3.73 (s, 3H), 3.98 (s, 3H), 5.60 (d, 1H), 6.63 (s, 1H), 6.68 (s, 1H)                  | 376 | [2-(2-methoxy-4,6-dimethylphenyl)-4-methoxy-6-fluoromethyl pyrimidin-5-yl] dipropylamine    |
| 15  | 2-methoxy-4,6-dimethylphenyl       | CHF <sub>2</sub>  | Me                                | 0.89 (t, 6H), 1.39 (m, 4H), 2.09 (s, 3H), 2.34 (s, 3H), 2.94 (t, 4H), 3.73 (s, 3H), 4.01 (s, 3H), 6.64 (s, 1H), 6.68 (s, 1H), 7.25 (t, 1H)                  | 394 | [2-(2-methoxy-4,6-dimethylphenyl)-4-methoxy-6-difluoromethyl pyrimidin-5-yl] dipropylamine  |

|    |                              |  |    |   |     |  |
|----|------------------------------|--|----|---|-----|--|
| 16 | 2-methoxy-4,6-dimethylphenyl | -CH(OH)CH <sub>3</sub>   | Me | 0.87 (t, 6H), 1.27 (d, 3H), 1.38 (m, 4H), 2.18 (s, 3H), 2.34 (s, 3H), 2.70 (dd, 1H), 2.94 (t, 4H), 3.34 (dd, 1H), 3.75 (s, 3H), 3.96 (s, 3H), 4.15 (m, 1H), 5.70 (br s, 1H), 6.64 (s, 1H), 6.69 (s, 1H)                   | 402 | 1-[5-(dipropylamino)-6-methoxy-2-(2-methoxy-4,6-dimethylphenyl)-pyrimidin-4-yl]-ethan-1-ol                   |
| 17 | 2-methoxy-4,6-dimethylphenyl | -C(CH <sub>3</sub> ) <sub>2</sub> OH   | Me | 0.88 (t, 6H), 1.23 (s, 6H), 1.37 (m, 4H), 2.17 (s, 3H), 2.34 (s, 3H), 2.90 (t, 4H), 3.10 (s, 2H), 3.73 (s, 3H), 3.96 (s, 3H), 6.61 (s, 1H), 6.63 (s, 1H), 6.68 (s, 1H)  | 416 | 1-[5-(dipropylamino)-6-methoxy-2-(2-methoxy-4,6-dimethylphenyl)-pyrimidin-4-yl]-propan-2-ol                  |
| 18 | 2-methoxy-4,6-dimethylphenyl |   | Me | 0.3-0.6 (m, 4H), 0.89 (t, 6H), 1.35 (t, 3H), 1.41 (m, 4H), 2.07 (s, 3H), 2.33 (s, 3H), 2.91 (t, 4H), 3.20 (ddd, 1H), 3.64 (ddd, 1H), 3.68 (s, 3H), 3.95 (s, 3H), 4.40 (q, 2H), 4.50 (ddt, 1H), 6.64 (s, 1H), 6.69 (s, 1H) | 430 | [4-(2-Cyclopropyl-2-fluoro-ethyl)-6-methoxy-2-(2-methoxy-4,6-dimethylphenyl)-pyrimidin-5-yl]-dipropyl-amine  |
| 19 | 2-methoxy-4,6-dimethylphenyl |  | Me | 0.2-0.6 (m, 4H), 0.88 (t, 6H), 1.00 (m, 1H), 1.41 (m, 4H), 2.17 (s, 3H), 2.34 (s, 3H), 2.91 (t, 4H), 2.92 (m, 2H), 3.27 (t, 1H), 3.46 (d, 1H), 3.74 (s, 3H), 3.96 (s, 3H), 5.75 (br s, 1H), 6.64 (s, 1H), 6.69 (s, 1H)    | 498 | [4-(2-Cyclopropyl-2-hydroxy-ethyl)-6-methoxy-2-(2-methoxy-4,6-dimethylphenyl)-pyrimidin-5-yl]-dipropyl-amine |



### Example 21

(Cyclopropylmethyl)[4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]propylamine [Formula I: Ar=6-methoxy-2,4-dimethylphenyl; R<sub>1</sub>=OCH<sub>3</sub>;

R<sub>2</sub>=N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>—); R<sub>3</sub>=CH<sub>3</sub>]

5 A. To a stirred solution of 4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidine-5-ylamine (900 mg, 3.3 mmol) in ethyl acetate (15 mL) is added triethylamine (430 mg, 4.25 mmol) and then cyclopropanecarbonyl chloride (416 mg, 4.0 mmol). The solution is stirred at room temperature for 1 h and then saturated aqueous sodium bicarbonate is added. The layers are separated and the aqueous layer further extracted with ethyl acetate. The combined  
10 organics are washed with water, brine, dried (sodium sulfate), filtered and concentrated to give cyclopropyl-N-[4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]carboxamide (1.02 g): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 0.85 (br, 2H), 1.1 (m, 2H), 1.2 (m, 1H), 2.05 (s, 3H), 2.34 (s, 3H), 2.38 (s, 3H), 3.65 (s, 3H), 3.98 (s, 3H), 6.60 (s, 1H), 6.64 (s, 1H); MS (CI) 342.


15 B. To a stirred solution of cyclopropyl-N-[4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]carboxamide (115 mg, 0.33 mmol) and 1-iodopropane (85 mg, 0.5 mmol) in DMF (2 mL) is added 60% sodium hydride (40 mg, 1.0 mmol). The mixture is heated to 55°C with stirring for 2 hr, cooled to room temperature and partitioned between ethyl acetate  
20 and saturated aqueous sodium bicarbonate. The layers are separated, the aqueous layer further extracted with ethyl acetate and the combined organics washed with water, brine, dried (sodium sulfate), filtered and concentrated to give cyclopropyl-N-[4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]-N-propylcarboxamide. An analytically pure sample is prepared by preparative TLC: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 0.65 (m, 2H), 0.90 (t, 3H), 1.05  
25 (m, 3H), 1.2 (m, 1H), 1.58 (m, 2H), 2.08 (s, 3H), 2.35 (s, 3H), 2.45 (s, 3H), 3.6 (m, 2H), 3.75 (s, 3H), 3.98 (s, 3H), 6.65 (s, 1H), 6.72 (s, 1H); MS (CI) 384.

C. To a stirred solution of cyclopropyl-N-[4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]-N-propylcarboxamide (90 mg, 0.2 mmol) in toluene at room temperature  
30 is added diisobutylaluminum hydride 1 M solution in hexanes (0.6 mL, 0.6 mmol). After 1 h the solution is acidified with 1 M hydrochloric acid, neutralized with 2 M sodium hydroxide solution

and extracted into ethyl acetate. The combined extracts are washed with water, brine, dried (sodium sulfate), filtered, concentrated, and chromatographed on silica (4:1 hexane-ether) to give 2-{5-[cyclopropylmethyl]propylamino}-4-methoxy-6-methylpyrimidin-2-yl}-3,5-dimethylphenol (61 mg): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ -0.02 (d, 2H), 0.38 (d, 2H), 0.78 (m, 1H), 0.85 (t, 3H), 1.28 (m, 2H), 2.28 (s, 3H), 2.58 (s, 3H), 2.78 (s, 3H), 2.82 (d, 2H), 2.95 (t, 2H), 4.02 (s, 3H), 6.60 (s, 1H), 6.72 (s, 1H); MS (CI) 356.

D. To 2-{5-[cyclopropylmethyl]propylamino}-4-methoxy-6-methylpyrimidin-2-yl}-3,5-dimethylphenol (35 mg, 0.1 mmol) in DMF (1 mL) is added cesium carbonate (163 mg, 0.5 mmol) and methyl iodide (0.1 mmol). The mixture is heated to 55 °C with stirring for 2 h, cooled to room temperature and partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The layers are separated, the aqueous layer further extracted with ethyl acetate and the combined organics washed with water, brine, dried (sodium sulfate), filtered and concentrated. The residue is purified by preparative TLC to give the title compound (22 mg): MS (CI) 370.


The EX#s 22-25 in the Table III may be prepared following the methods described in Example 21

| Table III   |    |         |   |
|---|----|---------|---|
|  |    |         |   |
| EX#   | R  | MS (CI) | NAME  |
| 22  | Et | 384     | Cycloproplmethyl-[2-(2-ethoxy-4,6-dimethylphenyl)-4-methoxy-6-methyl pyrimidin-5-yl] propyl-amine |

|    |                                     |     |  |
|----|-------------------------------------|-----|--|
| 23 | Pr                                  | 398 | Cyclopropylmethyl[2-(2-propoxy-4,6-dimethylphenyl)-4-methoxy-6-methylpyrimidin-5-yl] dipropylamine       |
| 24 | iPr                                 | 398 | Cyclopropylmethyl[2-(2-isopropoxy-4,6-dimethylphenyl)-4-methoxy-6-methylpyrimidin-5-yl] dipropylamine    |
| 25 | CH <sub>2</sub> CH <sub>2</sub> OMe | 414 | Cyclopropylmethyl[2-(2-ethoxymethoxy-4,6-dimethylphenyl)-4-methoxy-6-methylpyrimidin-5-yl] dipropylamine |

### Example 26

**[2-(dimethylamino)ethyl](cyclopropylmethyl)[6-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-4-methylpyrimidin-5-yl]amine**





[Formula I: Ar=6-methoxy-2,4-dimethylphenyl; R<sub>1</sub>=CH<sub>3</sub>; R<sub>2</sub>=N(CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>) (CH<sub>2</sub>—3=OCH<sub>3</sub>]

A. To a stirred solution of cyclopropyl-N-[4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-yl]carboxamide (115 mg, 0.33 mmol) and 2-(dimethylamino)ethyl chloride hydrochloride (72 mg, 0.5 mmol) in DMF (2 mL) is added 60% sodium hydride (40 mg, 1.0 mmol). The mixture is heated to 55°C with stirring for 2 h, cooled to room temperature and partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The layers are separated, the aqueous layer further extracted with ethyl acetate and the combined organics washed with water, brine, dried (sodium sulfate), filtered and concentrated to give N-[2-(dimethylamino)ethyl]cyclopropyl-N-[6-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-4-methylpyrimidin-5-yl]carboxamide (121 mg): MS (CI) 413.

B. To a stirred solution of N-[2-(dimethylamino)ethyl] cyclopropyl-N-[6-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-4-methylpyrimidin-5-yl]carboxamide (82 mg, 0.2 mmol) in

toluene at room temperature is added diisobutylaluminum hydride 1 M solution in hexanes (0.4 mL, 0.4 mmol). After 1 h the solution is acidified with 1 M hydrochloric acid, neutralized with 2 M sodium hydroxide solution and extracted into ethyl acetate. The combined extracts are washed with water, brine, dried (sodium sulfate), filtered, concentrated and purified by preparative TLC to give the title compound (61 mg): MS (CI) 399.

The EX#s 27-30 in the Table IV may be prepared following the methods described in Example 26.

| <div style="text-align: center;"> <b>Table IV</b><br/>  </div> |   |         |   |
|---|---|---------|---|
| EX#   | R   | MS (CI) | NAME  |
| 27  |  | 425     | Cyclopropylmethyl-[4-methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-yl]-(2-pyrrolidin-1-yl-ethyl)-amine |
| 28  |  | 442     | Cyclopropylmethyl-[4-methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-yl]-(2-morpholin-1-yl-ethyl)-amine  |
| 29  | OMe   | 400     | Cyclopropylmethyl-(2-methoxy-ethyl)-[4-methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-yl]-amine         |
| 30  |  | 440     | Cyclopropylmethyl-[4-methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-yl]-(2-piperidin-1-yl-ethyl)-amine  |



### Example 31


(Ethylpropyl)[6-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-4-methylpyrimidin-5-yl]amine

[Formula I: Ar=6-methoxy-2,4-dimethylphenyl; R<sub>1</sub>=CH<sub>3</sub>; R<sub>2</sub>=N(CH<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>);

R<sub>3</sub>=OCH<sub>3</sub>]

- 5           A suspension of 4-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-6-methylpyrimidin-5-ylamine (100 mg, 0.37 mmol), KI (166 mg, 1.0 mmol), K<sub>2</sub>CO<sub>3</sub> (138 mg, 1.0 mmol) and 2-bromopentane (151 mg, 1.0 mmol) in anhydrous acetonitrile (5 mL) in a pressure tube is heated at 100°C for 48 h. After cooling down, the mixture is partitioned between ethyl ether (50 mL) and brine (30 mL). The organic phase is washed with brine (30 mL), dried (MgSO<sub>4</sub>) and the
- 10 solvent removed under reduced pressure. Flash chromatography (25% ethyl acetate in hexanes) yields 6.9 mg of (ethylpropyl)[6-methoxy-2-(6-methoxy-2,4-dimethylphenyl)-4-methylpyrimidin-5-yl]amine. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 0.95 (t, 6H), 1.49 (m, 4H), 2.04 (s, 3H), 2.32 (s, 3H), 2.46 (s, 3H), 3.27 (m, 1H), 3.71 (s, 3H), 3.95 (s, 3H), 6.62 (s, 1H), 6.67 (s, 1H); MS (CI): 344.


**Table V**



| EX# | R <sub>1</sub>      | <sup>1</sup> H NMR (CDCl <sub>3</sub> )  | MS (CI) | NAME  |
|-----|---------------------|--|---------|---|
| 32  | 1-Propyl            | 1.00 (t, 3H), 1.59 (m, 2H), 2.06 (s, 3H), 2.33 (s, 3H), 2.49 (s, 3H), 3.11 (t, 2H), 3.72 (s, 3H), 3.96 (s, 3H), 6.62 (s, 1H), 6.67 (s, 1H).              | 368     | [4-Methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-yl]-propyl-amine      |
| 33  | Cyclopentyl         | 1.45 (m, 2H), 1.62 (m, 2H), 1.76 (m, 2H), 1.88 (m, 2H), 2.05 (s, 3H), 2.32 (s, 3H), 2.49 (s, 3H), 3.72 (s, 3H), 3.95 (s, 3H), 6.62 (s, 1H), 6.67 (2, 1H) | 342     | [4-Methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-yl]-cyclopentyl-amine |
| 34  | -CH <sub>2</sub> CN | 2.06 (s, 3H), 2.34 (s, 3H), 2.52 (s, 3H), 3.64 (t, 1H), 3.73 (s, 3H), 4.01 (s, 3H), 4.13 (d, 2H), 6.63 (s, 1H), 6.69 (s, 1H)                             | 313     | [4-Methoxy-2-(2-methoxy-4,6-dimethyl-phenyl)-6-methyl-pyrimidin-5-ylamino]-acetonitrile |

The EX#s 32-34 in the Table V may be prepared following the methods described in Example 31.


### Example 35

[2-(2-{5-[(cyclopropylmethyl)propylamino]-4-methoxy-6-methylpyrimidin-2-yl}-3,5-dimethylphenoxy)ethyl]dimethylamine [Formula I: Ar=2-(2-dimethylamineethoxy)-3,5-dimethylphenyl; R<sub>1</sub>=OCH<sub>3</sub>; R<sub>2</sub>=N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>—); R<sub>3</sub>=CH<sub>3</sub>]

- 5 A. To a stirred solution of 2-{5-[(cyclopropylmethyl)propylamino]-4-methoxy-6-methylpyrimidin-2-yl}-3,5-dimethylphenol (1.0 g, 2.8 mmol) in DMF (50 mL) is added cesium carbonate (3.26 g, 10.0 mmol) and 2-iodo-1-(1,1,2,2-tetramethyl-1-silapropoxy)ethane (1.43 g, 5.0 mmol). The mixture is heated to 80 °C with stirring for 2 h, cooled to room temperature and treated with methanol (40 mL) and 2 M hydrochloric acid (60 mL). The mixture is stirred overnight, basified with 4 M sodium hydroxide, extracted into ethyl acetate, washed with water, brine, dried (sodium sulfate), filtered and concentrated to give the crude 2-(2-{5-[(cyclopropylmethyl)propylamino]-4-methoxy-6-methylpyrimidin-2-yl}-3,5-dimethylphenoxy)ethan-1-ol (910 mg)
- 10
- 15 B. To the crude 2-(2-{5-[(cyclopropylmethyl)propylamino]-4-methoxy-6-methylpyrimidin-2-yl}-3,5-dimethylphenoxy)ethan-1-ol (40 mg, 0.1 mmol) in dichloromethane (1 mL) is added triethylamine (22 mg, 0.2 mmol) and methanesulfonyl chloride (22 mg, 0.4 mmol). After 1 hr the volatiles are removed by evaporation and the residue redissolved in acetonitrile (1 mL) and potassium carbonate (27 mg, 0.2 mmol) added followed by 1 M dimethylamine in THF (0.2 mL, 0.2 mmol). The mixture is heated to 60 °C with stirring for 2 h, cooled to room temperature partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The layers are separated, the aqueous layer further extracted with ethyl acetate and the combined organics washed with water, brine, dried (sodium sulfate), filtered and concentrated. The residue is purified by
- 20
- 25 preparative TLC to give the title compound (36 mg): MS (CI) 427.

The EX#s 36-52 in the Table VI may be prepared following the methods described in Example 35.

Table VI



| EX# | R <sub>7</sub> | MS (CI) | NAME   |
|-----|----------------|---------|--|
| 36  |                | 454     | {2-[2,4-Dimethyl-6-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropyl-amine            |
| 37  |                | 468     | {2-[2,4-Dimethyl-6-(2-piperidin-1-yl-ethoxy)-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropylamine              |
| 38  |                | 470     | {2-[2,4-Dimethyl-6-(2-morpholin-1-yl-ethoxy)-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropyl-amine             |
| 39  |                | 486     | {2-[2,4-Dimethyl-6-(2-thiomorpholin-4-yl-ethoxy)-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropyl-amine         |
| 40  |                | 482     | (2-{2,4-Dimethyl-6-[2-(4-methyl-piperidin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine  |
| 41  |                | 484     | (2-{2,4-Dimethyl-6-[2-(3-hydroxy-piperidin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine |
| 42  |                | 483     | (2-{2,4-Dimethyl-6-[2-(4-methyl-piperizin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine  |
| 43  |                | 456     | {2-[2-(2-Diethylamino-ethoxy)-4,6-dimethyl-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropyl-amine               |
| 44  |                | 558     | (2-{2-[2-(4-Benzyl-piperidin-1-yl)-ethoxy]-4,6-dimethyl-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine  |

|    |  |     |  |
|----|--|-----|--|
| 45 |  | 510 | (2-{2-[2-(Cyclohexyl-ethyl-amino)-ethoxy]-4,6-dimethyl-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine   |
| 46 |  | 482 | (2-{2,4-Dimethyl-6-[2-(2-methyl-piperidin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine  |
| 47 |  | 482 | (2-{2,4-Dimethyl-6-[2-(3-methyl-piperidin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine  |
| 48 |  | 482 | {2-[2-(2-Azepan-1-yl-ethoxy)-4,6-dimethyl-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropyl-amine                |
| 49 |  | 484 | (2-{2,4-Dimethyl-6-[2-(4-hydroxy-piperidin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine |
| 50 |  | 504 | (2-{2-[2-(Benzyl-methyl-amino)-ethoxy]-4,6-dimethyl-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine      |
| 51 |  | 522 | (2-{2,4-Dimethyl-6-[2-(octahydro-quinolin-1-yl)-ethoxy]-phenyl}-4-methoxy-6-methyl-pyrimidin-5-yl)-dipropyl-amine  |
| 52 |  | 451 | {2-[2-(2-Imidazol-1-yl-ethoxy)-4,6-dimethyl-phenyl]-4-methoxy-6-methyl-pyrimidin-5-yl}-dipropyl-amine              |

### Example 53

*Tert*-butyl-N-[4-methoxy-2-(2,6-dimethoxyphenyl)-6-methylpyrimidin-5-yl]carboxamide may be prepared following the method described in Step A of Example 521 starting from 4-methoxy-2-(2,6-dimethoxyphenyl)-6-methylpyrimidin-5-yl amine.


### Example 54

*Tert*-butyl-N-[n-propyl]-N-[4-methoxy-2-(2,6-dimethoxyphenyl)-6-methylpyrimidin-5-yl]carboxamide may be prepared following the method described in

Step B of Example 21 starting from *tert*-butyl-N-[4-methoxy-2-(2,6-dimethoxyphenyl)-6-methylpyrimidin-5-yl]carboxamide.

### Additional Examples

- 5 The following compounds can be prepared using the methods given in reaction schemes I - IV and further illustrated in the preceding examples.



| EX# | X                 | R                 | EX# | X               | R                 |
|-----|-------------------|-------------------|-----|-----------------|-------------------|
| 55  | OMe               | Me                | 80  | F               | Me                |
| 56  | OMe               | Et                | 81  | F               | Et                |
| 57  | OMe               | OCF <sub>3</sub>  | 82  | F               | OCF <sub>3</sub>  |
| 58  | OMe               | OCHF <sub>2</sub> | 83  | F               | OCHF <sub>2</sub> |
| 59  | OCF <sub>3</sub>  | Me                | 84  | Me              | Me                |
| 60  | OCF <sub>3</sub>  | Et                | 85  | Me              | Et                |
| 61  | OCF <sub>3</sub>  | OCF <sub>3</sub>  | 86  | Me              | OCF <sub>3</sub>  |
| 62  | OCF <sub>3</sub>  | OCHF <sub>2</sub> | 87  | Me              | OCHF <sub>2</sub> |
| 63  | OCHF <sub>2</sub> | Me                | 88  | Et              | Me                |
| 64  | OCHF <sub>2</sub> | Et                | 89  | Et              | Et                |
| 65  | OCHF <sub>2</sub> | OCF <sub>3</sub>  | 90  | Et              | OCF <sub>3</sub>  |
| 66  | OCHF <sub>2</sub> | OCHF <sub>2</sub> | 91  | Et              | OCHF <sub>2</sub> |
| 67  | Cl                | Me                | 92  | CF <sub>3</sub> | Me                |
| 68  | Cl                | Et                | 93  | CF <sub>3</sub> | Et                |
| 69  | Cl                | OCF <sub>3</sub>  | 94  | CF <sub>3</sub> | OCF <sub>3</sub>  |
| 70  | Cl                | OCHF <sub>2</sub> | 95  | CF <sub>3</sub> | OCHF <sub>2</sub> |

### 10 Example 96

#### Assay for CRF Receptor Binding Activity

As discussed above, the following assay is defined herein as a standard *in vitro* CRF receptor binding assay.

The pharmaceutical utility of compounds of this invention is indicated by the following assay for CRF1 receptor activity. The CRF receptor binding is performed using a modified version of the assay described by Grigoriadis and De Souza (*Methods in Neurosciences*, Vol. 5, 1991). IMR-32 human neuroblastoma cells, a cell-line that naturally expresses the CRF1 receptor, are grown to confluency in DMEM containing FBS.

To prepare receptor containing membranes cells are homogenized in wash buffer (50 mM Tris HCl, 10 mM MgCl<sub>2</sub>, 2 mM EGTA, pH 7.4) and centrifuged at 48,000 x g for 10 minutes at 4°C. The pellet is re-suspended in wash buffer and the homogenization and centrifugation steps are performed two additional times.

Membrane pellets containing CRF receptors are re-suspended in 50 mM Tris buffer pH 7.7 containing 10 mM MgCl<sub>2</sub> and 2 mM EDTA and centrifuged for 10 minutes at 48000g. Membranes are washed again and brought to a final concentration of 1500 mg/ml in binding buffer (Tris buffer above with 0.1 % BSA, 15 mM bacitracin and 0.01 mg/ml aprotinin.). For the binding assay, 100 ml of the membrane preparation are added to 96 well microtube plates containing 100 ml of <sup>125</sup>I-CRF (SA 2200 Ci/mmol, final concentration of 100 pM) and 50 ml of test compound. Binding is carried out at room temperature for 2 hours. Plates are then harvested on a Brandel 96 well cell harvester and filters are counted for gamma emissions on a Wallac 1205 Betaplate liquid scintillation counter. Non specific binding is defined by 1 mM cold CRF. IC<sub>50</sub> values are calculated with the non-linear curve fitting program RS/1 (BBN Software Products Corp., Cambridge, MA). The binding affinity for the compounds of Formula I expressed as IC<sub>50</sub> value, generally ranges from about 0.5 nanomolar to about 10 micromolar. Preferred compounds of Formula I exhibit IC<sub>50</sub> values of less than or equal to 1.5 micromolar, more preferred compounds of Formula I exhibit IC<sub>50</sub> values of less than 500 nanomolar, still more preferred compounds of Formula I exhibit IC<sub>50</sub> values of less than 100 nanomolar, and most preferred compound of Formula I exhibit IC<sub>50</sub> values of less than 10 nanomolar. The compounds shown in Examples 1-54 have been tested in this assay and found to exhibit IC<sub>50</sub> values of less than or equal to 4 micromolar.

### Example 97

#### Preparation of radiolabeled probe compounds of the invention

The compounds of the invention are prepared as radiolabeled probes by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope.

- 5 The radioisotope is preferably selected from at least one of carbon (preferably  $^{14}\text{C}$ ), hydrogen (preferably  $^3\text{H}$ ), sulfur (preferably  $^{35}\text{S}$ ), or iodine (preferably  $^{125}\text{I}$ ). Such radiolabeled probes are conveniently synthesized by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds. Such suppliers include Amersham Corporation, Arlington Heights, IL; Cambridge Isotope Laboratories, Inc.
- 10 Andover, MA; SRI International, Menlo Park, CA; Wizard Laboratories, West Sacramento, CA; ChemSyn Laboratories, Lexena, KS; American Radiolabeled Chemicals, Inc., St. Louis, MO; and Moravek Biochemicals Inc., Brea, CA.

- 15 Tritium labeled probe compounds are also conveniently prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas. Such preparations are also conveniently carried out as a custom radiolabeling by any of the suppliers listed in the preceding paragraph using the compound of the invention as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange
- 20 with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as appropriate.

### Example 98

#### Receptor autoradiography

- 25 Receptor autoradiography (receptor mapping) is carried out in vitro as described by Kuhar in sections 8.1.1 to 8.1.9 of Current Protocols in Pharmacology (1998) John Wiley & Sons, New York, using radiolabeled compounds of the invention prepared as described in the preceding Examples.

### 30 Example 99

#### Additional Aspects of Preferred Compounds of the Invention



The most preferred compounds of the invention are suitable for pharmaceutical use in treating human patients. Accordingly, such preferred compounds are non-toxic. They do not exhibit single or multiple dose acute or long-term toxicity, mutagenicity (e.g., as determined in a bacterial reverse mutation assay such as an Ames test),  
5     teratogenicity, tumorigenicity, or the like, and rarely trigger adverse effects (side effects) when administered at therapeutically effective dosages.

Preferably, administration of such preferred compounds of the invention at certain doses (e.g., doses yielding therapeutically effective in vivo concentrations or preferably doses of 10, 50, 100, 150, or 200 mg/kg -- preferably 150 mg/kg -- administered  
10     parenterally or preferably orally) does not result in prolongation of heart QT intervals (i.e., as determined by electrocardiography, e.g., in guinea pigs, minipigs or dogs). When administered daily for 5 or preferably ten days, such doses of such preferred compounds also do not cause liver enlargement resulting in an increase of liver to body weight ratio of more than 100%, preferably not more than 75% and more preferably not more than  
15     50% over matched controls in laboratory rodents (e.g., mice or rats). In another aspect such doses of such preferred compounds also preferably do not cause liver enlargement resulting in an increase of liver to body weight ratio of more than 50%, preferably preferably not more than 25%, and more preferably not more than 10% over matched untreated controls in dogs or other non-rodent animals.

20     In yet another aspect such doses of such preferred compounds also preferably do not promote the release of liver enzymes (e.g., ALT, LDH, or AST) from hepatocytes in vivo. Preferably such doses do not elevate such enzymes by more than 100%, preferably not by more than 75% and more preferably not by more than 50% over matched untreated controls in laboratory rodents. Similarly, concentrations (in culture media or  
25     other such solutions that are contacted and incubated with cells in vitro) equivalent to two, fold, preferably five-fold, and most preferably ten-fold the minimum in vivo therapeutic concentration do not cause release of any of such liver enzymes from hepatocytes in vitro.

Because side effects are often due to undesirable receptor activation or  
30     antagonism, preferred compounds of the invention exert their receptor-modulatory effects with high selectivity. This means that they do not bind to certain other receptors (i.e.,

other than CRF receptors) with high affinity, but rather only bind to, activate, or inhibit the activity of such other receptors with affinity constants of greater than 100 nanomolar, preferably greater than 1 micromolar, more preferably greater than 10 micromolar and most preferably greater than 100 micromolar. Such receptors preferably are selected from the group including ion channel receptors, including sodium ion channel receptors, neurotransmitter receptors such as alpha- and beta-adrenergic receptors, muscarinic receptors (particularly m1, m2, and m3 receptors), dopamine receptors, and metabotropic glutamate receptors; and also include histamine receptors and cytokine receptors, e.g., interleukin receptors, particularly IL-8 receptors. The group of other receptors to which preferred compounds do not bind with high affinity also includes GABA<sub>A</sub> receptors, bioactive peptide receptors (including NPY and VIP receptors), neurokinin receptors, bradykinin receptors (e.g., BK1 receptors and BK2 receptors), and hormone receptors (including thyrotropin releasing hormone receptors and melanocyte-concentrating hormone receptors).

#### **Example 99a**

##### **Absence of Sodium Ion Channel Activity**

Preferred compounds of the invention do not exhibit activity as Sodium ion channel blockers. Sodium channel activity may be measured a standard *in vitro* sodium channel binding assays such as the assay given by Brown et al. (J. Neurosci. (1986) 265: 17995-18004). Preferred compounds of the invention exhibit less than 15 percent inhibition, and more preferably less than 10 percent inhibition, of sodium channel specific ligand binding when present at a concentration of 4 uM. The sodium ion channel specific ligand used may be labeled batrachotoxinin, tetrodotoxin, or saxitoxin. Such assays, including the assay of Brown referred to above, are performed as a commercial service by CEREP, INC., Redmond, WA.

Alternatively, sodium ion channel activity may be measured *in vivo* in an assay of anti-epileptic activity. Anti-epileptic activity of compounds may be measured by the ability of the compounds to inhibit hind limb extension in the supra maximal electro shock model. Male Han Wistar rats (150-200mg) are dosed i.p. with a suspension of 1 to 20 mg of test compound in 0.25% methylcellulose 2 hr. prior to test. A visual

observation is carried out just prior to testing for the presence of ataxia. Using auricular electrodes a current of 200 mA, duration 200 millisec, is applied and the presence or absence of hind limb extension is noted. Preferred compounds of the invention do not exhibit significant anti-epileptic activity at the  $p < 0.1$  level of significance or more preferably at the  $p < 0.05$  level of significance as measured using a standard parametric assay of statistical significance such as a student's T test.

### **Example 99b**

#### **Optimal in vitro half-life**

- 10 Compound half-life values ( $t_{1/2}$  values) may be determined via the following standard liver microsomal half-life assay. Liver microsomes obtained from pooled liver samples and prepared so that the P-450 enzyme content is approximately 0.5 nmol/ mg protein. Reactions are preformed in a 5ml well deep-well plate as follows:
- Phosphate buffer: 19 mL 0.1 M  $\text{NaH}_2\text{PO}_4$ , 81 mL 0.1  $\text{Na}_2\text{HPO}_4$ , pH 7.4 with  $\text{H}_3\text{PO}_4$ .
- 15 CoFactor Mixture: 16.2 mg NADP, 45.4 mg Glucose-6-phosphate in 4 mL 100 mM  $\text{MgCl}_2$ . Glucose-6-phosphate dehydrogenase: 214.3 ul glucose-6-phosphate dehydrogenase, 1285.7 ul distilled water
- Starting Reaction Mixture: 3 mL CoFactor Mixture, 1.2 mL Glucose-6-phosphate dehydrogenase 6 identical samples wells each containing 25 ul microsomes, 5 ul of test
- 20 compound (from a 100 uM stock), and 399 ul 0.1 M phosphate buffer, pH 7.4, are prepared. A seventh well containing 25 ul microsomes, 399 ul 0.1 M phosphate buffer, pH 7.4, and 5 ul (from a 100 uM stock) of a compound, e.g. DIAZEPAM, CLOZEPINE, with known metabolic properties is used as a positive control. Reactions are preincubated at 39 °C for 10 minutes. 71 ul Starting Reaction Mixture is added to 5 of the
- 25 6 reaction wells and to the positive control well, 71 ul 100 mM  $\text{MgCl}_2$  is added to the sixth reaction well, which is used as a negative control. At each time point (0, 1, 3, 5, and 10 minutes) 75 ul reaction is pipetted into a 96-well deep-well plate reaction well containing 75 ul ice-cold acetonitrile. Samples are vortexed and centrifuged 10 minutes at 6000 rpm (Sorval T 6000D rotor). Supernatant, 75 ul from each reaction well, is
- 30 transferred to a 96-well plate containing 150 ul internal standard per well. The remaining test compound is quantitated via LCMS and Compound concentration vs time is plotted

and commercially available statistical software is used to extrapolate to the  $t_{1/2}$  value of the test compound.

Preferred compounds of the invention exhibit in vitro  $t_{1/2}$  values of greater than 10 minutes and less than 4 hours. Most preferred compounds of the invention exhibit in vitro  $t_{1/2}$  values of between 30 minutes and 1 hour in human liver microsomes.

### **Example 99c**

#### **MDCK Toxicity**

Toxicity of a test compound may be assessed by measuring the effect of the compound on ATP production by MDCK cells.

MDCK cells, product no. CCL-34, are purchased from ATCC, Manassas, VA and maintained in sterile conditions by the methods described in the supplier's production information sheet. The PACKARD BIOSCIENCE, (Groningen, The Netherlands) ATP-LITE-M Luminescent ATP detection kit, product no. 6016941, may be used to monitor ATP production in MDCK cells.

Prior to assay 1  $\mu$ l of test compound or control sample is pipetted into PACKARD (Meriden, CT) clear bottom 96-well plates. Test compounds and control samples are diluted in DMSO to give final concentration in the assay of 10 micromolar, 100 micromolar, or 200 micromolar. Control samples are drug compounds having known toxicity properties.

Confluent MDCK cells are trypsinized, harvested, and diluted to a concentration of  $0.1 \times 10^6$  cells/ml with warm ATCC Eagle's Minimum Essential Medium (catalog # 30-2003). Warm Eagle's MEM without cells (100ul) is pipetted in five wells a 96-well plate. These wells are used to determine the standard curve. Cells in Eagle's MEM (100ul or 10,000 cells) are pipetted into the remaining wells of the 96-well plates. All samples are incubated at 37 °C under carbogen (95% O<sub>2</sub>, 5% CO<sub>2</sub>) for 2 hours with constant shaking. After incubation 50 ul mammalian cell lysis solution is added to well of the 96-well plates, wells are covered with Packard topseal stickers, and plates are shaken at approximately 700 rpm on a microtiter shaker for 2 minutes.

During the incubation, Packard ATP Lite-M reagents are allowed to equilibrate to room temperature. Once equilibrated the lyophilized substrate solution is reconstituted in

5.5 mls of substrate buffer solution (from kit). Lyophilized ATP standard solution is reconstituted in deionized water to give a 10 mM stock. Standard (10 ul), diluted so that a 10 ul aliquot yields a final concentration of 200 nM, 100 nM, 50 nM, 25 nM, or 12.5 nM is added to each of the five standard curve wells, which do not contain cells.

- 5 Substrate solution (50 ul) is added to all wells. Wells are covered with Packard topseal stickers, and plates are shaken at approximately 700 rpm on a microtiter shaker for 2 minutes. A white Packard sticker is attached to the bottom of each plate and samples are dark adapted by wrapping plates in foil and placing in the dark for 10 minutes. Luminescence is then quantitated at 22 °C using a luminescence counter, e.g.
- 10 Packard TopCount Microplate Scintillation and Luminescence Counter or Tecan Spectrafluor plus.

- Luminescence values at each concentration are compared to the values computed from the standard curve for that concentration. Preferred test compounds exhibit luminescence values 80 % or more of the standard, or preferably 90 % or more of the standard, when 10 micromolar concentration of the test compound is used. When a 100 micromolar concentration of the test compound is used, preferred test compounds exhibit luminescence values 50 % or more of the standard, or more preferably 80 % or more of the standard.
- 15

- The disclosures of all articles and references mentioned in this application,
- 20 including patents, are incorporated herein by reference.

- The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that
- 25 modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.